

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 September 2001 (07.09.2001)


PCT

(10) International Publication Number
WO 01/64654 A1

- (51) International Patent Classification⁷: C07D 239/48, 401/12, 417/12, 413/12, A61K 31/505, 31/506
- (21) International Application Number: PCT/GB01/00782
- (22) International Filing Date: 26 February 2001 (26.02.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0004888.4 1 March 2000 (01.03.2000) GB
- (71) Applicant (for all designated States except MG, US): ASTRAZENECA AB [SE/SE]; S-151 85 Sodertalje (SE).
- (71) Applicant (for MG only): ASTRAZENECA UK LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB).
- (72) Inventors; and
(75) Inventors/Applicants (for US only): PEASE, Elizabeth, Janet [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). BREAU, Gloria, Anne [US/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). MORRIS, Jeffrey, James [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).
- (74) Agent: BRYANT, Tracey; AstraZeneca, Global Intellectual Property, P.O. Box 272, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4GR (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


[Continued on next page]

(54) Title: PYRIMIDINE COMPOUNDS




(I)

(57) Abstract: Pyrimidine derivatives of formula (I) wherein: Q₁ and Q₂ are independently selected from aryl or carbon linked heteroaryl optionally substituted as defined within; and one of Q₁ and Q₂ or both Q₁ and Q₂ is substituted on a ring carbon by one group selected from sulphonamoyl, N-(C₁₋₄alkyl)sulphonamoyl (optionally substituted by halo or hydroxy), N,N-di-(C₁₋₄alkyl)sulphonamoyl (optionally substituted by halo or hydroxy), C₁₋₄alkylsulphonyl (optionally substituted by halo or hydroxy) or a substituent of the formula (Ia) or (Ia'): wherein Q₁, Q₂, G, R¹, Y, Z, Q₃, n and m are as defined within; and pharmaceutically acceptable salts and *in vivo* hydrolysable esters thereof are described. Processes for their manufacture, pharmaceutical compositions and their use as cyclin-dependent serine/threonine kinase (CDK) inhibitors are also described.



(Ia)



(Ia')



(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

PYRIMIDINE COMPOUNDS

The invention relates to pyrimidine derivatives, or pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof, which possess cell-cycle inhibitory activity and are accordingly useful for their anti cell proliferation (such as anti-cancer) activity and are therefore useful in methods of treatment of a warm-blooded animal, such as man. The invention also relates to processes for the manufacture of said pyrimidine derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments or use in the production of an anti-cell-proliferation effect in a warm-blooded animal such as man.

A family of intracellular proteins called cyclins play a central role in the cell cycle. The synthesis and degradation of cyclins is tightly controlled such that their level of expression fluctuates during the cell cycle. Cyclins bind to cyclin-dependent serine/threonine kinases (CDKs) and this association is essential for CDK (such as CDK1, CDK2, CDK4 and/or CDK6) activity within the cell. Although the precise details of how each of these factors combine to regulate CDK activity is poorly understood, the balance between the two dictates whether or not the cell will progress through the cell cycle.

The recent convergence of oncogene and tumour suppresser gene research has identified regulation of entry into the cell cycle as a key control point of mitogenesis in tumours. Moreover, CDKs appear to be downstream of a number of oncogene signalling pathways. Disregulation of CDK activity by upregulation of cyclins and/or deletion of endogenous inhibitors appears to be an important axis between mitogenic signalling pathways and proliferation of tumour cells.


Accordingly it has been recognised that an inhibitor of cell cycle kinases, particularly inhibitors of CDK2, CDK4 and/or CDK6 (which operate at the S-phase, G1-S and G1-S phase respectively) should be of value as a selective inhibitor of cell proliferation, such as growth of mammalian cancer cells.

The present invention is based on the discovery that certain 2,4-pyrimidine compounds surprisingly inhibit the effects of cell cycle kinases showing selectivity for CDK2, CDK4 and CDK6, and thus possess anti-cell-proliferation properties. Such properties are expected to be of value in the treatment of disease states associated with aberrant cell cycles and cell

proliferation such as cancers (solid tumours and leukemias), fibroproliferative and differentiative disorders, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, atherosclerosis, arterial restenosis, autoimmune diseases, acute and chronic inflammation, bone diseases and ocular diseases with retinal vessel proliferation.

According to the invention there is provided a pyrimidine derivative of the formula

(I):




(I)

10 wherein:

Q_1 and Q_2 are independently selected from aryl or carbon linked heteroaryl; and one of Q_1 and Q_2 or both Q_1 and Q_2 is substituted on a ring carbon by one group selected from sulphamoyl, *N*-(C_{1-4} alkyl)sulphamoyl (optionally substituted by halo or hydroxy), *N,N*-di-(C_{1-4} alkyl)sulphamoyl (optionally substituted by halo or hydroxy), C_{1-4} alkylsulphonyl

15 (optionally substituted by halo or hydroxy) or a substituent of the formula (Ia) or (Ia'):



wherein:

Y is -NHS(O)₂-, -S(O)₂NH- or -S(O)₂-;

20 Z is R^aO-, R^bR^cN-, R^dS-, R^eR^fNNR^g-, C₃₋₈cycloalkyl, phenyl or a heterocyclic group; wherein said phenyl, C₃₋₈cycloalkyl or heterocyclic group are optionally substituted on a ring carbon by one or more groups selected from R^h; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from Rⁱ;

R^a , R^b , R^c , R^d , R^e , R^f and R^g are independently selected from hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, phenyl, heterocyclic group and C_{3-8} cycloalkyl; wherein said C_{1-4} alkyl, C_{2-4} alkenyl and C_{3-8} cycloalkyl are optionally substituted by one or more groups selected from R^i ;

n is 0 or 1;

- 5 m is 1, 2 or 3, in addition m may be 0 when Z is C_{3-8} cycloalkyl, phenyl or a heterocyclic group;

Q_3 is a nitrogen linked heterocycle; wherein said heterocycle is optionally substituted on a ring carbon by one or more groups selected from R^k ; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group

- 10 selected from R^m

G is -O-, -S- or -NR²-;

R^2 is selected from hydrogen, C_{1-6} alkyl, C_{3-6} alkenyl and C_{3-6} alkynyl; wherein said C_{1-6} alkyl, C_{3-6} alkenyl and C_{3-6} alkynyl are optionally substituted by one or more groups selected from R^n ;

- 15 R^1 is selected from hydrogen, halo, hydroxy, nitro, amino, N -(C_{1-3} alkyl)amino, N,N -di-(C_{1-3} alkyl)amino, cyano, trifluoromethyl, trichloromethyl, C_{1-3} alkyl [optionally substituted by 1 or 2 substituents independently selected from halo, cyano, amino, N -(C_{1-3} alkyl)amino, N,N -di-(C_{1-3} alkyl)amino, hydroxy and trifluoromethyl], C_{3-5} alkenyl [optionally substituted by up to three halo substituents, or by one trifluoromethyl substituent],
20 C_{3-5} alkynyl, C_{1-3} alkoxy, mercapto, C_{1-3} alkylsulphanyl, carboxy and C_{1-3} alkoxycarbonyl;

Q_1 is optionally substituted on a ring carbon by one to four substituents independently selected from halo, mercapto, nitro, formyl, formamido, carboxy, cyano, amino, ureido, carbamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl [wherein said C_{1-4} alkyl, C_{2-4} alkenyl and C_{2-4} alkynyl are optionally substituted by one or more groups selected from R^o], C_{1-4} alkanoyl,
25 C_{1-4} alkoxycarbonyl, heterocyclic group, C_{1-4} alkylS(O)_a wherein a is 0 or 1 [optionally substituted by hydroxy], N' -(C_{1-4} alkyl)ureido, N',N' -di-(C_{1-4} alkyl)ureido, N' -(C_{1-4} alkyl)- N' -(C_{1-4} alkyl)ureido, N - C_{1-4} alkylamino, N,N -di-(C_{1-4} alkyl)amino, N - C_{1-4} alkylcarbamoyl, N,N -di-(C_{1-4} alkyl)carbamoyl and C_{1-4} alkanoylamino;

- 30 and also independently, or in addition to, the above substituents, Q_1 may be optionally substituted by one to two substituents independently selected from aryl, C_{3-8} cycloalkyl and a heterocyclic group; wherein said aryl, C_{3-8} cycloalkyl or heterocyclic group may be optionally

substituted on a ring carbon by one or more groups selected from R^p; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^q;

and also independently, or in addition to, the above substituents, Q₁ may be optionally substituted by one C₁₋₄alkoxy or by one hydroxy substituent;

Q₂ is optionally substituted on a ring carbon by one to four substituents independently selected from halo, hydroxy, mercapto, nitro, formyl, formamido, carboxy, cyano, amino, ureido, carbamoyl, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy [wherein said C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl and C₁₋₄alkoxy are optionally substituted by one or more groups selected from R^j], C₁₋₄alkanoyl, C₁₋₄alkoxycarbonyl, heterocyclic group, C₁₋₄alkylS(O)_a, wherein a is 0 or 1 [optionally substituted by hydroxy], N¹-(C₁₋₄alkyl)ureido, N¹,N¹-di-(C₁₋₄alkyl)ureido, N¹-(C₁₋₄alkyl)-N-(C₁₋₄alkyl)ureido, N¹,N¹-di-(C₁₋₄alkyl)-N-(C₁₋₄alkyl)ureido, N-C₁₋₄alkylamino, N,N-di-(C₁₋₄alkyl)amino, N-C₁₋₄alkylcarbamoyl, N,N-di-(C₁₋₄alkyl)carbamoyl, C₂₋₄alkenyloxy, C₂₋₄alkynyloxy and C₁₋₄alkanoylamino;

and also independently, or in addition to, the above substituents, Q₂ may be optionally substituted by one to two substituents independently selected from aryl, C₃₋₈cycloalkyl or a heterocyclic group; wherein said aryl, C₃₋₈cycloalkyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R^s; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^t;

R^j, Rⁿ, R^o and R^r are independently selected from hydroxy, halo, amino, cyano, formyl, formamido, carboxy, nitro, mercapto, carbamoyl, sulphamoyl, N-C₁₋₄alkylamino, N,N-di-(C₁₋₄alkyl)amino, C₁₋₄alkanoyl, C₁₋₄alkanoyloxy, C₁₋₄alkoxy, C₁₋₄alkoxycarbonyl, N-C₁₋₄alkylcarbamoyl, N,N-di-(C₁₋₄alkyl)carbamoyl, C₁₋₄alkanoylamino, C₁₋₄alkylS(O)_a, wherein a is 0 to 2, C₁₋₄alkylsulphonylamino, N-(C₁₋₄alkyl)sulphamoyl, N-(C₁₋₄alkyl)₂sulphamoyl, N-(C₁₋₄alkyl)carbamoyl, N-(C₁₋₄alkyl)₂carbamoyl, phenyl, phenylthio, phenoxy, C₃₋₈cycloalkyl and a heterocyclic group; wherein said phenyl, phenylthio, phenoxy, C₃₋₈cycloalkyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R^u; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^v;

R^h , R^k , R^p , R^s and R^u are independently selected from hydroxy, halo, amino, cyano, formyl, formamido, carboxy, nitro, mercapto, carbamoyl, sulphamoyl, C_{1-4} alkyl [optionally substituted by one or more groups selected from halo, cyano, amino, N - C_{1-4} alkylamino, N,N -di- $(C_{1-4}$ alkyl)amino or hydroxy], C_{2-4} alkenyl [optionally substituted by one or more groups selected from halo], C_{2-4} alkynyl, N - C_{1-4} alkylamino, N,N -di- $(C_{1-4}$ alkyl)amino, C_{1-4} alkanoyl, C_{1-4} alkanoyloxy, C_{1-4} alkoxy [optionally substituted by one or more groups selected from halo], C_{1-4} alkoxycarbonyl, N - C_{1-4} alkylcarbamoyl, N,N -di- $(C_{1-4}$ alkyl)carbamoyl, C_{1-4} alkanoylamino, C_{1-4} alkylS(O)_a wherein a is 0 to 2, C_{1-4} alkylsulphonylamino, N -(C_{1-4} alkyl)sulphamoyl, N -(C_{1-4} alkyl)₂sulphamoyl, phenyl, C_{3-8} cycloalkyl and a heterocyclic group; and

R^i , R^q , R^t and R^v are independently selected from C_{1-4} alkyl, C_{1-4} alkanoyl, C_{1-4} alkylsulphonyl, C_{1-4} alkoxycarbonyl, carbamoyl, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl; or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

“Aryl” is a fully or partially unsaturated, mono or bicyclic carbon ring that contains 4-12 atoms. Preferably “aryl” is a monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms. More preferably “aryl” is phenyl, naphthyl, tetralinyl or indanyl. Particularly “aryl” is phenyl, naphthyl or indanyl. More particularly “aryl” is phenyl.

A “carbon linked heteroaryl” is a fully unsaturated, 5- or 6- membered monocyclic ring or 9- or 10- membered bicyclic ring of which at least one atom is chosen from nitrogen, sulphur or oxygen. This ring is linked via a carbon atom to the -NH- (for Q_1) or G (for Q_2). Preferably “carbon linked heteroaryl” is furanyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyrazolyl, furazanyl, triazolyl, thiadiazolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazinyl, indolyl, quinolyl or benzimidazolyl. More preferably “carbon linked heteroaryl” is pyridyl, thiazolyl or pyrazolyl. Particularly “carbon linked heteroaryl” is pyridyl.

A “heterocyclic group” is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 4-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH₂- group can optionally be replaced by a -C(O)-, and a ring sulphur atom may be optionally oxidised to form S-oxide(s). Preferably a “heterocyclic group” is pyrrolidinyl, morpholino, piperidyl, pyridyl, pyranyl, pyrrolyl, isothiazolyl, indolyl, quinolyl, thienyl, furyl,

1,3-benzodioxolyl, thiadiazolyl, piperazinyl, isoxazolyl, thiazolyl, thiazolidinyl, pyrrolidinyl, thiomorpholino, pyrazolyl, pyrrolinyl, homopiperazinyl, tetrahydropyranyl, imidazolyl, pyrimidyl, pyrazinyl, pyridazinyl, isoxazolyl, 4-pyridone, 1-isoquinolone, 2-pyrrolidone, 4-thiazolidone, imidazo[1,2-a]pyridine or 3-aza-8-oxabicyclo[3,2,1]hexane. More preferably a

5 “heterocyclic group” is pyrrolidinyl, morpholino, piperidyl, isoxazolyl, thiadiazolyl, thiazolyl, pyridyl, indolyl, thienyl, furyl, piperazinyl, thiomorpholino, pyrazolyl, imidazolyl, 2-pyrrolidone, imidazo[1,2-a]pyridine or 3-aza-8-oxabicyclo[3,2,1]hexane. Particularly a “heterocyclic group” is morpholino, isoxazolyl, thiadiazolyl, thiazolyl or pyridyl.

A “nitrogen linked heterocycle” is a saturated, partially saturated or fully unsaturated,

10 mono or bicyclic ring containing 4-12 atoms, one atom of which is a nitrogen atom (attached to form an amide as shown) and the other atoms are either all carbon atoms or they are carbon atoms and 1-3 heteroatoms chosen from nitrogen, sulphur or oxygen, wherein a -CH₂- group can optionally be replaced by a -C(O)- and a ring sulphur atom may be optionally oxidised to form the S-oxides. It will be appreciated that in forming this nitrogen link, the nitrogen atom

15 is not quaternised, i.e. a neutral compound is formed. Preferably “nitrogen linked heterocycle” is pyrrol-1-yl, pyrrolin-1-yl, pyrrolidin-1-yl, imidazol-1-yl, imidazolin-1-yl, imidazolidin-1-yl, pyrazol-1-yl, pyrazolin-1-yl, pyrazolidin-1-yl, triazol-1-yl, piperidin-1-yl, piperazin-1-yl, morpholino, thiomorpholino, indol-1-yl, indolidin-1-yl or benzimidazol-1-yl. More preferably “nitrogen linked heterocycle” is piperidin-1-yl.

20 In this specification the term “alkyl” includes both straight and branched chain alkyl groups but references to individual alkyl groups such as “propyl” are specific for the straight chain version only. An analogous convention applies to other generic terms. “Halo” is fluoro, chloro, bromo and iodo.

Examples of C₂₋₄alkenyl are vinyl and allyl; examples of C₂₋₆alkenyl are C₃₋₅alkenyl, vinyl and allyl; examples of C₃₋₆alkenyl are C₃₋₅alkynyl and allyl; an example of C₃₋₆alkynyl is propyn-2-yl; examples of C₂₋₄alkynyl are ethynyl and propyn-2-yl; examples of C₂₋₆alkynyl are ethynyl and propyn-2-yl; examples of C₁₋₄alkanoyl are acetyl and propionyl; examples of C₁₋₄alkoxycarbonyl are C₁₋₃alkoxycarbonyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl and *tert*-butoxycarbonyl; examples of C₁₋₄alkylene are methylene, ethylene

30 and propylene; examples of C₁₋₄alkyl are C₁₋₃alkyl, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl and *tert*-butyl; examples of C₁₋₆alkyl are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, *tert*-butyl and 3-methylbutyl; examples of C₁₋₄alkoxy are

C₁₋₃alkoxy, methoxy, ethoxy, propoxy, isopropoxy and butoxy; an example of C₂₋₄alkenyloxy is allyloxy; an example of C₂₋₄alkynyloxy is propynyloxy; examples of C₁₋₄alkylS(O)_a,
 wherein a is 0 or 1 are C₁₋₃alkylsulphanyl, methylthio, ethylthio, propylthio, methylsulphanyl, ethylsulphanyl and propylsulphanyl; examples of C₁₋₄alkylS(O)_a, wherein a is 0 to 2 are
 5 methylthio, ethylthio, propylthio, methylsulphanyl, ethylsulphanyl, propylsulphanyl, mesyl, ethylsulphonyl and propylsulphonyl; examples of C₁₋₄alkylsulphonyl are mesyl and ethylsulphonyl; examples of N-C₁₋₄alkylcarbamoyl are N-methylcarbamoyl, N-ethylcarbamoyl and N-propylcarbamoyl; examples of N,N-di-(C₁₋₄alkyl)-carbamoyl are N,N-dimethylcarbamoyl, N-ethyl-N-methylcarbamoyl and N,N-diethylcarbamoyl; examples of
 10 N-C₁₋₄alkylamino are N-(C₁₋₃alkyl)amino, methylamino, ethylamino and propylamino; examples of N,N-di-(C₁₋₄alkyl)amino are N,N-di-(C₁₋₃alkyl)amino, dimethylamino, N-ethyl-N-methylamino, diethylamino, N-methyl-N-propylamino and dipropylamino; examples of C₁₋₄alkanoylamino are acetamido, propionamido and butyramido; examples of C₃₋₈cycloalkyl are cyclopropyl, cyclopentyl and cyclohexyl; examples of C₁₋₄alkanoyl are
 15 acetyl and propionyl; examples of C₁₋₄alkanoyloxy are acetyloxy and propionyloxy; examples of N'-(C₁₋₄alkyl)ureido are N'-methylureido and N'-ethylureido; examples of N',N'-di-(C₁₋₄alkyl)ureido are N',N'-dimethylureido, N',N'-diisopropylureido and N'-methyl-N'-propylureido; examples of N'-(C₁₋₄alkyl)-N-(C₁₋₄alkyl)ureido are N'-methyl-N-ethylureido and N'-methyl-N-methylureido; examples of
 20 N',N'-di-(C₁₋₄alkyl)-N-(C₁₋₄alkyl)ureido are N',N'-dimethyl-N-ethylureido and N'-methyl-N'-propyl-N-butylureido; examples of N-(C₁₋₄alkyl)sulphamoyl are N-methylsulphamoyl and N-isopropylsulphamoyl; examples of N,N-di-(C₁₋₄alkyl)sulphamoyl are N-methyl-N-ethylsulphamoyl and N,N-dipropylsulphamoyl; and examples of C₁₋₄alkylsulphonylamino are mesylamino,
 25 ethylsulphonylamino and propylsulphonylamino.

A suitable pharmaceutically acceptable salt of a pyrimidine derivative of the invention is, for example, an acid-addition salt of a pyrimidine derivative of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic,
 30 citric or maleic acid. In addition a suitable pharmaceutically acceptable salt of a pyrimidine derivative of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium

salt, an ammonium salt or a salt with an organic base which affords a physiologically acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

The compounds of the formula (I) may be administered in the form of a pro-drug
5 which is broken down in the human or animal body to give a compound of the formula (I).
Examples of pro-drugs include *in vivo* hydrolysable esters of a compound of the formula (I).

An *in vivo* hydrolysable ester of a compound of the formula (I) containing carboxy or hydroxy group is, for example, a pharmaceutically acceptable ester which is hydrolysed in the human or animal body to produce the parent acid or alcohol. Suitable pharmaceutically
10 acceptable esters for carboxy include C₁₋₆alkoxymethyl esters for example methoxymethyl, C₁₋₆alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, C₃₋₈cycloalkoxycarbonyloxyC₁₋₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl and may be
15 formed at any carboxy group in the compounds of this invention.

An *in vivo* hydrolysable ester of a compound of the formula (I) containing a hydroxy group includes inorganic esters such as phosphate esters and α -acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and
20 2,2-dimethylpropionyloxy-methoxy. A selection of *in vivo* hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and *N*-(dialkylaminoethyl)-*N*-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl. Examples of substituents on benzoyl include morpholino and piperazino linked
25 from a ring nitrogen atom via a methylene group to the 3- or 4- position of the benzoyl ring.

Some compounds of the formula (I) may have chiral centres and/or geometric isomeric centres (E- and Z- isomers), and it is to be understood that the invention encompasses all such optical, diastereo-isomers and geometric isomers that possess CDK inhibitory activity.

30 The invention relates to any and all tautomeric forms of the compounds of the formula (I) that possess CDK inhibitory activity.

It is also to be understood that certain compounds of the formula (I) can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which possess CDK inhibitory activity.

- 5 Particular preferred compounds of the invention comprise a pyrimidine derivative of the formula (I), or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, wherein R¹, Q₁, Q₂, and G have any of the meanings defined hereinbefore, or any of the following values. Such values may be used where appropriate with any of the definitions, claims or embodiments defined hereinbefore or hereinafter.
- 10 Preferably Q₁ and Q₂ are independently selected from phenyl and pyridyl.
Preferably Q₁ is phenyl.
Preferably Q₂ is phenyl or pyridyl.
Preferably Q₁ is phenyl and Q₂ is selected from phenyl or pyridyl.
More preferably Q₁ and Q₂ are phenyl.
- 15 Preferably one of Q₁ and Q₂ or both Q₁ and Q₂ is substituted on a ring carbon by one group selected from sulphamoyl, *N*-(C₁₋₄alkyl)sulphamoyl, *N,N*-di-(C₁₋₄alkyl)sulphamoyl (optionally substituted by hydroxy), C₁₋₄alkylsulphonyl or a substituent of formula (Ia) wherein:
- Y is -S(O)₂NH- or -S(O)₂-;
- 20 Z is R^aO-, R^bR^cN- or a heterocyclic group; wherein heterocyclic group are optionally substituted on a ring carbon by one or more groups selected from R^h;
- R^a, R^b and R^c are independently selected from hydrogen, C₁₋₄alkyl and phenyl;
- n is 0;
- m is 2 or in addition m may be 0 when Z is a heterocyclic group.
- 25 More preferably one of Q₁ and Q₂ or both of Q₁ and Q₂ is substituted on a ring carbon by one group selected from sulphamoyl, *N*-(C₁₋₄alkyl)sulphamoyl, *N,N*-di-(C₁₋₄alkyl)sulphamoyl (optionally substituted by hydroxy), C₁₋₄alkylsulphonyl or a substituent of formula (Ia) wherein:
- Y is -S(O)₂NH- or -S(O)₂-;
- 30 Z is R^aO-, R^bR^cN-, thiazolyl, isoxazolyl, thiadiazolyl, pyridyl or morpholino; wherein thiazolyl, isoxazolyl, thiadiazolyl, pyridyl or morpholino are optionally substituted on a ring carbon by one or more methyl;

R^a , R^b and R^c are independently selected from hydrogen, C_{1-4} alkyl and phenyl;

n is 0;

m is 2 or in addition m may be 0 when Z is thiazolyl, isoxazolyl, thiadiazolyl or pyridyl.

- 5 Particularly one of Q_1 and Q_2 or both of Q_1 and Q_2 is substituted on a ring carbon by one group selected from sulphamoyl, mesyl, *N*-(2-diethylaminoethyl)sulphamoyl, 2-(*N*-methyl-*N*-phenylamino)ethylsulphonyl, 2-morpholinoethylsulphonyl, *N*-(5-methylthiadiazol-2-yl)sulphamoyl, *N,N*-di-(2-hydroxyethyl)sulphamoyl, *N*-(thiazol-2-yl)sulphamoyl, *N*-(3,4-dimethylisoxazol-5-yl)sulphamoyl,
- 10 *N*-(pyrid-2-yl)sulphamoyl and *N*-methylsulphamoyl.

More particularly one of Q_1 and Q_2 or both of Q_1 and Q_2 is substituted on a ring carbon by one group selected from sulphamoyl and *N*-methylsulphamoyl.

Preferably it is Q_1 that is substituted by one group selected from sulphamoyl, *N*-(C_{1-4} alkyl)sulphamoyl (optionally substituted by halo or hydroxy),

- 15 *N,N*-di-(C_{1-4} alkyl)sulphamoyl (optionally substituted by halo or hydroxy), C_{1-4} alkylsulphonyl (optionally substituted by halo or hydroxy) or a substituent of the formula (Ia) or (Ia'); and Q_2 is optionally additionally substituted by one group selected from sulphamoyl.

More preferably it is Q_1 that is substituted in the para- or meta- position relative to the -NH- by sulphamoyl, *N*-(C_{1-4} alkyl)sulphamoyl (optionally substituted by halo or hydroxy),

- 20 *N,N*-di-(C_{1-4} alkyl)sulphamoyl (optionally substituted by halo or hydroxy), C_{1-4} alkylsulphonyl (optionally substituted by halo or hydroxy) or a substituent of the formula (Ia) or (Ia'); and Q_2 is optionally additionally substituted by one group selected from sulphamoyl para to G.

Particularly it is Q_1 that is substituted in the para- position relative to the -NH- by sulphamoyl, *N*-(C_{1-4} alkyl)sulphamoyl (optionally substituted by halo or hydroxy),

- 25 *N,N*-di-(C_{1-4} alkyl)sulphamoyl (optionally substituted by halo or hydroxy), C_{1-4} alkylsulphonyl (optionally substituted by halo or hydroxy) or a substituent of the formula (Ia) or (Ia'); and Q_2 is optionally additionally substituted by one group selected from sulphamoyl para to G.

In one aspect of the invention preferably G is -O-.

In another aspect of the invention preferably G is -S-.

- 30 In a further aspect of the invention preferably G is -NR²-.

In one aspect of the invention preferably G is -O- or -NR²-.

In one aspect of the invention when G is -NR²-, preferably R² is hydrogen.

In another aspect of the invention when G is -NR²-, preferably R² is not hydrogen.

Preferably G is -O- or -NR²- wherein R² is selected from hydrogen, C₁₋₆alkyl and C₃₋₆alkenyl; wherein said C₁₋₆alkyl and C₃₋₆alkenyl are optionally substituted by one or more halo, cyano or phenyl.

- 5 More preferably G is -O- or -NR²- wherein R² is selected from hydrogen, C₁₋₆alkyl and C₃₋₆alkenyl; wherein said C₁₋₆alkyl and C₃₋₆alkenyl are optionally substituted by one or more halo or phenyl.

Particularly G is -O-, -NH-, -(4,4,4-trifluorobutyl)N-, -(3-bromo-2-propenyl)N- or -(3-phenyl-2-propenyl)N-.

- 10 More particularly G is -O- or -NH-.

Preferably R¹ is hydrogen or halo.

More preferably R¹ is hydrogen, chloro or bromo.

- Preferably Q₁ is optionally substituted by one C₁₋₄alkoxy substituent and is substituted by one group selected from sulphamoyl, *N*-(C₁₋₄alkyl)sulphamoyl (optionally substituted by
15 halo or hydroxy), *N,N*-di-(C₁₋₄alkyl)sulphamoyl (optionally substituted by halo or hydroxy), C₁₋₄alkylsulphonyl (optionally substituted by halo or hydroxy) or a substituent of the formula (Ia) or (Ia').

More preferably Q₁ is substituted by one group selected from sulphamoyl, *N*-(C₁₋₄alkyl)sulphamoyl or *N,N*-di-(C₁₋₄alkyl)sulphamoyl.

- 20 Preferably Q₂ is unsubstituted or substituted by one or two groups selected from halo, cyano, C₁₋₄alkyl, C₁₋₄alkoxy and a heterocyclic group.

More preferably Q₂ is optionally substituted on a ring carbon by one to two substituents independently selected from halo, cyano, methyl, methoxy and morpholino.

- Particularly Q₂ is optionally substituted on a ring carbon by one to two substituents
25 independently selected from cyano and methoxy.

Preferably Q₂ is phenyl, 2-morpholinophenyl, 2-cyanophenyl, 4-bromophenyl, 2-fluoro-5-methylphenyl, 4-methoxyphenyl or 4-sulphamoylphenyl.

More preferably Q₂ is phenyl, 2-cyanophenyl, 4-methoxyphenyl or 4-sulphamoylphenyl.

- 30 Therefore, in a preferred aspect of the invention there is provided a pyrimidine derivative of the formula (I) as depicted above, wherein:

Q_1 and Q_2 are independently selected from phenyl and pyridyl; and one of Q_1 and Q_2 or both Q_1 and Q_2 is substituted on a ring carbon by one group selected from sulphamoyl, N -(C_{1-4} alkyl)sulphamoyl, N,N -di-(C_{1-4} alkyl)sulphamoyl (optionally substituted by hydroxy), C_{1-4} alkylsulphonyl or a substituent of formula (Ia) wherein:

5 Y is $-S(O)_2NH-$ or $-S(O)_2-$;

Z is R^aO- , R^bR^cN- or a heterocyclic group; wherein heterocyclic group are optionally substituted on a ring carbon by one or more groups selected from R^h ;

R^a , R^b and R^c are independently selected from hydrogen, C_{1-4} alkyl and phenyl;

n is 0;

10 m is 2 or in addition m may be 0 when Z is a heterocyclic group;

G is $-O-$ or $-NR^2-$ wherein R^2 is selected from hydrogen, C_{1-6} alkyl and C_{3-6} alkenyl; wherein said C_{1-6} alkyl and C_{3-6} alkenyl are optionally substituted by one or more halo or phenyl;

R^1 is hydrogen or halo;

15 or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

Therefore, in a more preferred aspect of the invention there is provided a pyrimidine derivative of the formula (I) as depicted above, wherein:

Q_1 is phenyl optionally substituted by one C_{1-4} alkoxy substituent and Q_2 is phenyl optionally substituted by one or more halo, cyano, methyl, methoxy and morpholino; and Q_1

20 is substituted in the para- position relative to the $-NH-$ by one group selected from sulphamoyl, mesyl, N -(2-diethylaminoethyl)sulphamoyl,

2-(N -methyl- N -phenylamino)ethylsulphonyl, 2-morpholinoethylsulphonyl,

N -(5-methylthiadiazol-2-yl)sulphamoyl, N,N -di-(2-hydroxyethyl)sulphamoyl,

N -(thiazol-2-yl)sulphamoyl, N -(3,4-dimethylisoxazol-5-yl)sulphamoyl,

25 N -(pyrid-2-yl)sulphamoyl and N -methylsulphamoyl; and Q_2 is optionally substituted by one group selected from sulphamoyl;

G is $-O-$, $-NH-$, $-(4,4,4\text{-trifluorobutyl})N-$, $-(3\text{-bromo-2-propenyl})N-$ or $-(3\text{-phenyl-2-propenyl})N-$;

R^1 is hydrogen, chloro or bromo

30 or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

In one aspect of the invention preferred compounds of the invention are those of Examples 1, 20, 21, 29 or 31 or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.


In a further aspect of the invention preferred compounds of the invention include any
5 one of the Examples or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

Preferred aspects of the invention are those which relate to the compound or a pharmaceutically acceptable salt thereof.


A pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, may be prepared by any process known to be applicable to
10 the preparation of chemically-related compounds. Such processes, when used to prepare a pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, are provided as a further feature of the invention and are illustrated by the following representative examples in which, unless otherwise stated R^1 , Q_1 , Q_2 and G have any of the meanings defined hereinbefore for a pyrimidine derivative of the formula (I)
15 and unless another substituent is drawn on ring Q_1 or Q_2 the ring may bear any of the substituents described hereinbefore (optionally protected as necessary). Where a substituent is drawn on ring Q_1 , this includes (unless stated otherwise) the possibilities of the substituent being on ring Q_2 in addition to, or instead of the substituent being on ring Q_1 . Necessary starting materials may be obtained by standard procedures of organic chemistry (see for
20 example, Advanced Organic Chemistry (Wiley-Interscience), Jerry March - also useful for general guidance on reaction conditions and reagents). The preparation of such starting materials is described within the accompanying non-limiting processes and Examples. Alternatively necessary starting materials are obtainable by analogous procedures to those illustrated which are within the ordinary skill of an organic chemist.

25 Thus, as a further feature of the invention there are provided the following processes which comprises of:-

a) for compounds of formula (I) where G is $-NR^2-$; reacting a pyrimidine of formula (II):




wherein L is a displaceable group as defined below, with a compound of formula (III):




where G is $-\text{NR}^2-$;

5 b) reaction of a pyrimidine of formula (IV):




wherein L is a displaceable group as defined below, with a compound of formula (V):




10


c) for compounds of formula (I) wherein the sidechain is of formula (Ia) and Y is $-\text{S}(\text{O})_2\text{NH}-$; by reaction of a compound of formula (VI):



15 where L is a displaceable group; with an amine of formula (VII):




d) for compounds of formula (I) wherein the sidechain is of formula (Ia) and Y is $-\text{NHS}(\text{O})_2-$ by reaction of an amine of formula (VIII):



20


(VIII)

with a compound of formula (IX):



5 where L is a displaceable group;

e) for compounds of formula (I) wherein the sidechain is of formula (Ia'); by reaction of a compound of formula (VI) with an amine of formula (X):



10 and thereafter if necessary:

- i) converting a compound of the formula (I) into another compound of the formula (I);
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

The skilled reader will also appreciate that process c) can also be use to make

15 compounds of formula (I) wherein one of Q₁ and Q₂ or both Q₁ and Q₂ is substituted on a ring carbon by one group selected from sulphamoyl, *N*-(C₁₋₄alkyl)sulphamoyl (optionally substituted by halo or hydroxy) or *N,N*-di-(C₁₋₄alkyl)sulphamoyl (optionally substituted by halo or hydroxy).

L is a displaceable group, suitable values for L are for example, a halo, sulphonyloxy or sulphur group, for example a chloro, bromo, methanesulphonyloxy, toluene-4-sulphonyloxy, mesyl, methylthio and methylsulphinyl.

Specific reaction conditions for the above reactions are as follows:-

Process a)


Pyrimidines of formula (II) and compounds of formula (III) may be reacted together:

25 i) optionally in the presence of a suitable acid, for example an inorganic acid such as hydrochloric acid or sulphuric acid, or an organic acid such as acetic acid or formic acid. The reaction is preferably carried out in a suitable inert solvent or diluent, for example dichloromethane (DCM), acetonitrile, butanol, tetramethylene sulphone, tetrahydrofuran, 1,2-dimethoxyethane, *N,N*-dimethylformamide, *N,N*-dimethylacetamide or

N-methylpyrrolidin-2-one, and at a temperature in the range, for example, 0° to 150°C, conveniently at or near reflux temperature; or

ii) under standard Buchwald conditions (for example see *J. Am. Chem. Soc.*, **118**, 7215; *J. Am. Chem. Soc.*, **119**, 8451; *J. Org. Chem.*, **62**, 1568 and 6066) for example in the presence of
 5 palladium acetate, in a suitable solvent for example an aromatic solvent such as toluene, benzene or xylene, with a suitable base for example an inorganic base such as caesium carbonate or an organic base such as potassium-*t*-butoxide, in the presence of a suitable ligand such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl and at a temperature in the range of 25 to 80°C.

10 Pyrimidines of the formula (II) may be prepared according to the following scheme:



wherein L is a displaceable group as defined above.

Compounds of formula (IIB) and (III) are commercially available or are prepared by processes known in the art.


15 Process b)

Pyrimidines of formula (IV) and anilines of formula (V) may be reacted together, i) in the presence of a suitable solvent for example a ketone such as acetone or an alcohol such as ethanol or butanol or an aromatic hydrocarbon such as toluene or *N*-methyl pyrrolidine, optionally in the presence of a suitable acid such as those defined above (or a suitable Lewis

20 acid) and at a temperature in the range of 0°C to reflux, preferably reflux; or

ii) under standard Buchwald conditions as described above.

Pyrimidines of formula (IV) are prepared according to the following scheme:



wherein L is a displaceable group as defined above.


The anilines of formula (V) are commercially available or are prepared by processes known in the art.

Pyrimidines of the formula (IVA) are commercially available or may be prepared by, for example, reacting a compound of formula (IVA) in which L is -OH (i.e. a uracil), with POCl₃ to give a compound of formula (IVA) in which L is -Cl.

Process c)

10 Compounds of formula (VI) and amines of formula (VII) may be coupled together in the presence of a base, for example a tertiary amine such as triethylamine and in the presence of a catalyst for example dimethylaminopyridine. Suitable solvents for the reaction include nitriles such as acetonitrile and amides such as dimethylformamide. The reaction is conveniently performed at a temperature in the range of from 0 to 120°C.

15 Compounds of formula (VI) (for example when L is chlorine) may be prepared according to the following scheme:




wherein Pg is a suitable sulphur protecting group such as those described below.

Amines of formula (VII) and amines of formula (VIa) are commercially available or are prepared by processes known in the art.

Process d)

Compounds of formula (IX) and amines of formula (VIII) may be coupled together under the conditions described in process c) above.

Amines of formula (VIII) may be prepared according to the following scheme:



5

wherein Pg is a suitable amino protecting group such as those described hereinbelow.

Compounds of formula (IX) are commercially available or are prepared by processes known in the art.

Process e)

10


Compounds of formula (VI) and amines of formula (X) may be coupled together under the conditions described in process c) above.

Amines of formula (X) are commercially available or are prepared by processes known in the art.

Examples of conversions of a compound of formula (I) into another compound of

15 formula (I) are:


i) where G is -NR²-; conversion of R² as hydrogen into other R² for example:



wherein L is a displaceable group;

ii) where G is -NR²-; conversion of R² as a substituted side chain into another substituted

20 side chain, for example:



wherein Ms is methanesulphonyl, and Nu is a nucleophile that introduces a substituent that is an optional substituent for R^2 as defined in formula (I) (NB the hydroxyl moiety does not necessarily have to be on the terminal carbon as depicted above);

- 5 iii) conversion of one side chain of formula (Ia) into another side chain of formula (Ia).
- iv) conversion of one value of R^1 into another value of R^1 , using standard techniques, for example, conversion of R^1 as hydroxy into C_{1-4} alkoxy.

The skilled reader will appreciate that the formation of the side chain (Ia) or (Ia') described in Processes c), d) and e) above and of the sidechain R^2 in i) and ii) above may also
 10 be performed on intermediates.

A preferred process of the invention is Process b).

It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately
 15 following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution
 20 reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group

using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halo group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating;
5 oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Conventional protecting groups may be used in accordance with standard
10 practice (for illustration see T.W. Green, Protective Groups in Organic Synthesis, John Wiley and Sons, 1991). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a
15 methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali
20 metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with
25 a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an
30 arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with

a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a thio group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acetyl or benzoyl group may be removed, for example, by cleavage with sodium and ammonia.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

Many of the intermediates defined herein are novel, for example, those of the formula II and IV and these are provided as a further feature of the invention.

ASSAYS

As stated hereinbefore the pyrimidine derivative defined in the present invention possesses anti-cell-proliferation activity such as anti-cancer activity which is believed to arise from the CDK inhibitory activity of the compound. These properties may be assessed, for example, using the procedure set out below:-

CDK4 Inhibition Assay

The following abbreviations have been used :-

HEPES is *N*-(2-Hydroxyethyl)piperazine-*N'*-(2-ethanesulfonic acid)

DTT is Dithiothreitol

PMSF is Phenylmethylsulfonyl fluoride

The compounds were tested in an *in vitro* kinase assay in 96 well format using Scintillation Proximity Assay (SPA - obtained from Amersham) for measuring incorporation of [γ -33-P]-Adenosine Triphosphate into a test substrate (GST-Retinoblastoma). In each well was placed the compound to be tested (diluted in DMSO and water to correct concentrations) and in control wells either p16 as an inhibitor control or DMSO as a positive control.

Approximately 0.5 μ l of CDK4/Cyclin D1 partially-purified enzyme (amount dependent on enzyme activity) diluted in 25 μ l incubation buffer was added to each well then 20 μ l of GST-Rb/ATP/ATP33 mixture (containing 0.5 μ g GST-Rb and 0.2 μ M ATP and 0.14 μ Ci [γ -33-P]-Adenosine Triphosphate), and the resulting mixture shaken gently, then incubated at room temperature for 60 minutes.

To each well was then added 150 μ L stop solution containing (0.8mg/well of Protein A-PVT SPA bead (Amersham)), 20pM/well of Anti-Glutathione Transferase, Rabbit IgG (obtained from Molecular Probes), 61mM EDTA and 50mM HEPES pH 7.5 containing 0.05% sodium azide.

The plates were sealed with Topseal-S plate sealers, left for two hours then spun at 2500rpm, 1124xg., for 5 minutes. The plates were read on a Topcount for 30 seconds per well.

The incubation buffer used to dilute the enzyme and substrate mixes contained 50mM HEPES pH7.5, 10mM MnCl₂, 1mM DTT, 100 μ M Sodium vanadate, 100 μ M NaF, 10mM Sodium Glycerophosphate, BSA (1mg/ml final).

As a control, another known inhibitor of CDK4 may be used in place of p16.

Test substrate

In this assay only part of the retinoblastoma (Science 1987 Mar13;235(4794):1394-1399; Lee W.H., Bookstein R., Hong F., Young L.J., Shew J.Y., Lee E.Y.) was used, fused to a GST tag. PCR of retinoblastoma amino acids 379-928 (obtained from retinoblastoma plasmid ATCC pLRbRNL) was performed, and the sequence cloned into pGEX 2T fusion vector (Smith D.B. and Johnson, K.S. Gene 67, 31 (1988); which contained a tac promoter for inducible expression, internal lac I^s gene for use in any E.Coli host, and a coding region for thrombin cleavage - obtained from Pharmacia Biotech) which was used to amplify amino acids 792-928. This sequence was again cloned into pGEX 2T.

The retinoblastoma 792-928 sequence so obtained was expressed in E.Coli (BL21 (DE3) pLysS cells) using standard inducible expression techniques, and purified as follows.

E.coli paste was resuspended in 10ml/g of NETN buffer (50mM Tris pH 7.5, 120mM NaCl, 1mM EDTA, 0.5%v/v NP-40, 1mM PMSF, 1ug/ml leupeptin, 1ug/ml aprotinin and 1ug/ml pepstatin) and sonicated for 2 x 45 seconds per 100ml homogenate. After centrifugation, the supernatant was loaded onto a 10ml glutathione Sepharose column (Pharmacia Biotech, Herts, UK), and washed with NETN buffer. After washing with kinase buffer (50mM HEPES pH 7.5, 10mM MgCl₂, 1mM DTT, 1mM PMSF, 1ug/ml leupeptin, 1ug/ml aprotinin and 1ug/ml pepstatin) the protein was eluted with 50mM reduced glutathione in kinase buffer. Fractions containing GST-Rb(792-927) were pooled and dialysed overnight against kinase buffer. The final product was analysed by Sodium Dodeca Sulfate (SDS) PAGE (Polyacrylamide gel) using 8-16% Tris-Glycine gels (Novex, San Diego, USA).

CDK4 and Cyclin D1

CDK4 and Cyclin D1 were cloned from RNA from MCF-7 cell line (obtained from ATCC number:HTB22, breast adenocarcinoma line) as follows. The RNA was prepared from MCF-7 cells, then reverse transcribed using oligo dT primers. PCR was used to amplify the complete coding sequence of each gene [CDK4 amino acids 1-303; Ref. Cell 1992 Oct 16; 71(2): 323-334; Matsushima H., Ewen M.E., Stron D.K., Kato J.Y., Hanks S.K., Roussel M.F., Sherr C.J. and Cyclin D1 amino acids 1-296; Ref. Cold Spring Harb. Symp. Quant. Biol., 1991; 56:93-97; Arnold A., Motokura T., Bloom T., Kronenburg, Ruderman J., Juppner H., Kim H.G.].

After sequencing the PCR products were cloned using standard techniques into the insect expression vector pVL1393 (obtained from Invitrogen 1995 catalogue number : V1392-20). The PCR products were then dually expressed [using a standard virus Baculogold co-infection technique] into the insect SF21 cell system (Spodoptera Frugiperda cells derived from ovarian tissue of the Fall Army Worm -Commercially available).

The following Example provides details of the production of Cyclin D1/CDK4 in SF21 cells (in TC100 + 10% FBS(TCS) + 0.2% Pluronic) having dual infection MOI 3 for each virus of Cyclin D1 & CDK4.

Example production of Cyclin D1/CDK4

SF21 cells grown in a roller bottle culture to 2.33×10^6 cells/ml were used to inoculate 10 x 500 ml roller bottles at 0.2×10^6 cells/ml. The roller bottles were incubated on a roller rig at 28°C.

After 3 days (72 hrs.) the cells were counted, and the average from 2 bottles found to be 1.86×10^6 cells/ml. (99% viable). The cultures were then infected with the dual viruses at an MOI 3 for each virus.

10 x 500ml were infected with JS303 Cyclin D1 virus titre - 9×10^7 pfu/ml. JS304
5 CDK4 virus titre - 1×10^8 pfu/ml.

Cyclin D1 $\frac{1.86 \times 10^6 \times 500 \times 3}{0.9 \times 10^8} = 31$ ml of virus for each 500 ml. bottle.

CDK4 $\frac{1.86 \times 10^6 \times 500 \times 3}{1 \times 10^8} = 28$ ml of virus for each 500 ml. bottle.

10 The viruses were mixed together before addition to the cultures, and the cultures returned to the roller rig 28°C.

After 3 days (72 hrs.) post infection the 5 Litres of culture was harvested. The total cell count at harvest was 1.58×10^6 cells/ml.(99% viable). The cells were spun out at 2500rpm, 30 mins., 4°C in Heraeus Omnifuge 2.0 RS in 250 ml lots. The supernatant was
15 discarded. 20 pellets of $\sim 4 \times 10^8$ cells/pellet were snap frozen in LN₂ and stored at -80°C in CCRF cold room. The SF21 cells were then hypotonically lysed by resuspending in lysis buffer (50mM HEPES pH 7.5, 10mM magnesium chloride, 1mM DTT, 10mM glycerophosphate, 0.1mM PMSF, 0.1mM sodium fluoride, 0.1mM sodium orthovanadate, 5ug/ml aprotinin, 5ug/ml leupeptin and 20% w/v sucrose), and adding ice cold deionised
20 water. After centrifugation, the supernatant was loaded onto a Poros HQ/M 1.4/100 anion exchange column (PE Biosystems, Hertford, UK). CDK4 and Cyclin D1 were coeluted with 375mM NaCl in lysis buffer, and their presence checked by western blot, using suitable anti-CDK4 and anti-Cyclin D1 antibodies (obtained from Santa Cruz Biotechnology, California, US).

25 p16 control (Nature 366, 704-707; 1993; Serrano M, Hannon GJ, Beach D)

p16 (the natural inhibitor of CDK4/Cyclin D1) was amplified from HeLa cDNA (Hela cells obtained from ATCC CCL2, human epitheloid carcinoma from cervix; Cancer Res. 12: 264, 1952), cloned into pTB 375 NBSE which had a 5' His tag, and transformed using standard techniques into BL21 (DE3) pLysS cells (obtained from Promega; Ref. Studier F.W. and Moffat B.A., J. Mol. Biol., 189, 113, 1986). A 1 litre culture was grown to the appropriate
30 OD then induced with IPTG to express p16 overnight. The cells were then lysed by sonication in 50mM sodium phosphate, 0.5M sodium chloride, PMSF, 0.5µg/ml leupeptin and 0.5µg/ml

aprotinin. The mixture was spun down, the supernatant added to nickel chelate beads and mixed for 1 ½ hours. The beads were washed in sodium phosphate, NaCl pH 6.0 and p16 product eluted in sodium phosphate, NaCl pH 7.4 with 200mM imidazole.

The pTB NBSE was constructed from pTB 375 NBPE as follows :-

5 pTB375

The background vector used for generation of pTB 375 was pZEN0042 (see UK patent 2253852) and contained the tetA/tetR inducible tetracycline resistance sequence from plasmid RP4 and the cer stability sequence from plasmid pKS492 in a pAT153 derived background. pTB375 was generated by the addition of an expression cassette consisting of the T7 gene 10 promoter, multiple cloning site and T7 gene 10 termination sequence. In addition, a terminator sequence designed to reduce transcriptional readthrough from the background vector was included upstream of the expression cassette.

pTB 375 NBPE

The unique EcoRI restriction site present in pTB 375 was removed. A new multiple 15 cloning site containing the recognition sequences for the restriction enzymes NdeI, BamHI, PstI and EcoRI was introduced into pTB 375 between the NdeI and BamHI sites destroying the original BamHI site present in pTB 375.

pTB 375 NBSE

A new multiple cloning site containing the recognition sequences for the restriction 20 enzymes NdeI, BamHI, SmaI and EcoRI was introduced into pTB 375 NBPE between the NdeI and EcoRI sites. The oligonucleotide containing these restriction sites also contained 6 histidine codons located between the NdeI and BamHI sites in the same reading frame as the initiator codon (ATG) present within the NdeI site.

By analogy to the above, assays designed to assess inhibition of CDK2 and CDK6 25 may be constructed. CDK2 (EMBL Accession No. X62071) may be used together with Cyclin A or Cyclin E (see EMBL Accession No. M73812), and further details for such assays are contained in PCT International Publication No. WO99/21845, the relevant Biochemical & Biological Evaluation sections of which are hereby incorporated by reference.

If using CDK2 with Cyclin E partial co-purification may be achieved as follows:-
30 Sf21 cells are resuspended in lysis buffer (50mM Tris pH 8.2, 10mM MgCl₂, 1mM DTT, 10mM glycerophosphate, 0.1mM sodium orthovanadate, 0.1mM NaF, 1mM PMSF, 1ug/ml leupeptin and 1ug/ml aprotinin) and homogenised for 2 minutes in a 10ml Dounce

homogeniser. After centrifugation, the supernatant is loaded onto a Poros HQ/M 1.4/100 anion exchange column (PE Biosystems, Hertford, UK). CDK2 and Cyclin E are coeluted at the beginning of a 0-1M NaCl gradient (run in lysis buffer minus protease inhibitors) over 20 column volumes. Co-elution is checked by western blot using both anti-CDK2 and
5 anti-Cyclin E antibodies (Santa Cruz Biotechnology, California, US).

Although the pharmacological properties of the compounds of the formula (I) vary with structural change, in general activity possessed by compounds of the formula (I) in the above assays may be demonstrated at IC_{50} concentrations or doses in the range 250 μ M to 1nM.

10 When tested in the above *in vitro* assay the CDK2 inhibitory activity of Example 23 was measured as $IC_{50} = 0.347\mu$ M.

The *in vivo* activity of the compounds of the present invention may be assessed by standard techniques, for example by measuring inhibition of cell growth and assessing cytotoxicity.

15 Inhibition of cell growth may be measured by staining cells with Sulforhodamine B (SRB), a fluorescent dye that stains proteins and therefore gives an estimation of amount of protein (i.e. cells) in a well (see Boyd, M. R. (1989) Status of the NCI preclinical antitumour drug discovery screen. *Prin. Prac Oncol* 10:1-12). Thus, the following details are provided of measuring inhibition of cell growth:-

20 Cells were plated in appropriate medium in a volume of 100 μ l in 96 well plates; media was Dulbecco's Modified Eagle media for MCF-7, SK-UT-1B and SK-UT-1. The cells were allowed to attach overnight, then inhibitor compounds were added at various concentrations in a maximum concentration of 1% DMSO (v/v). A control plate was assayed to give a value for cells before dosing. Cells were incubated at 37°C, (5% CO₂) for three days.

25 At the end of three days TCA was added to the plates to a final concentration of 16% (v/v). Plates were then incubated at 4°C for 1 hour, the supernatant removed and the plates washed in tap water. After drying, 100 μ l SRB dye (0.4% SRB in 1% acetic acid) was added for 30 minutes at 37°C. Excess SRB was removed and the plates washed in 1% acetic acid. The SRB bound to protein was solubilised in 10mM Tris pH7.5 and shaken for 30 minutes at
30 room temperature. The ODs were read at 540nm, and the concentration of inhibitor causing 50% inhibition of growth was determined from a semi-log plot of inhibitor concentration versus absorbance. The concentration of compound that reduced the optical density to below

that obtained when the cells were plated at the start of the experiment gave the value for toxicity.

Typical IC_{50} values for compounds of the invention when tested in the SRB assay are in the range 1mM to 1nM.

5 According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in association with a pharmaceutically acceptable diluent or carrier.

10 The composition may be in a form suitable for oral administration, for example as a tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) as a sterile solution, suspension or emulsion, for topical administration as an ointment or cream or for rectal administration as a suppository.

In general the above compositions may be prepared in a conventional manner using conventional excipients.

15 The pyrimidine will normally be administered to a warm-blooded animal at a unit dose within the range 5-5000 mg per square meter body area of the animal, i.e. approximately 0.1-100 mg/kg, and this normally provides a therapeutically-effective dose. A unit dose form such as a tablet or capsule will usually contain, for example 1-250 mg of active ingredient. Preferably a daily dose in the range of 1-50 mg/kg is employed. However the daily dose will
20 necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

25 According to a further aspect of the present invention there is provided a pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man.

30 We have found that the pyrimidine derivatives defined in the present invention, or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, are effective cell cycle inhibitors (anti-cell proliferation agents), which property (without being bound by theory) is believed to arise from their CDK inhibitory properties. Accordingly the compounds of the present invention are expected to be useful in the treatment of diseases or medical conditions mediated alone or in part by CDK enzymes, i.e. the compounds may be used to produce a

CDK inhibitory effect in a warm-blooded animal in need of such treatment. Thus the compounds of the present invention provide a method for treating the proliferation of malignant cells characterised by inhibition of CDK enzymes, i.e. the compounds may be used to produce an anti-proliferative effect mediated alone or in part by the inhibition of CDKs.

5 Such a pyrimidine derivative of the invention is expected to possess a wide range of anti-cancer properties as CDKs have been implicated in many common human cancers such as leukaemia and breast, lung, colon, rectal, stomach, prostate, bladder, pancreas and ovarian cancer. Thus it is expected that a pyrimidine derivative of the invention will possess anti-cancer activity against these cancers. It is in addition expected that a pyrimidine
10 derivative of the present invention will possess activity against a range of leukaemias, lymphoid malignancies and solid tumours such as carcinomas and sarcomas in tissues such as the liver, kidney, prostate and pancreas. In particular such compounds of the invention are expected to slow advantageously the growth of primary and recurrent solid tumours of, for example, the colon, breast, prostate, lungs and skin. More particularly such compounds of the
15 invention, or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, are expected to inhibit the growth of those primary and recurrent solid tumours which are associated with CDK, especially those tumours which are significantly dependent on CDK for their growth and spread, including for example, certain tumours of the colon, breast, prostate, lung, vulva and skin.

20 It is further expected that a pyrimidine derivative of the present invention will possess activity against other cell-proliferation diseases in a wide range of other disease states including leukemias, fibroproliferative and differentiative disorders, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, atherosclerosis, arterial restenosis, autoimmune diseases, acute and chronic inflammation,
25 bone diseases and ocular diseases with retinal vessel proliferation.

Thus according to this aspect of the invention there is provided a pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore for use as a medicament; and the use of a pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as
30 defined hereinbefore in the manufacture of a medicament for use in the production of an anti-cancer, cell cycle inhibitory (anti-cell-proliferation) effect in a warm-blooded animal such

as man. Particularly, a cell cycle inhibitory effect is produced at the S or G1-S phase by inhibition of CDK2, CDK4 and/or CDK6, especially CDK2.

According to a further feature of this aspect of the invention there is provided a method for producing an anti-cancer, cell cycle inhibitory (anti-cell-proliferation) effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a pyrimidine derivative as defined immediately above. Particularly, an inhibitory effect is produced at the S or G1-S phase by inhibition of CDK2, CDK4 and/or CDK6, especially CDK2.

As stated above the size of the dose required for the therapeutic or prophylactic treatment of a particular cell-proliferation disease will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. A unit dose in the range, for example, 1-100 mg/kg, preferably 1-50 mg/kg is envisaged.

The CDK inhibitory activity defined hereinbefore may be applied as a sole therapy or may involve, in addition to a compound of the invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. In the field of medical oncology it is normal practice to use a combination of different forms of treatment to treat each patient with cancer. In medical oncology the other component(s) of such conjoint treatment in addition to the cell cycle inhibitory treatment defined hereinbefore may be: surgery, radiotherapy or chemotherapy. Such chemotherapy may cover three main categories of therapeutic agent:

- (i) other cell cycle inhibitory agents that work by the same or different mechanisms from those defined hereinbefore;
- (ii) cytostatic agents such as antioestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene, idoxifene), progestogens (for example megestrol acetate), aromatase inhibitors (for example anastrozole, letrozole, vorazole, exemestane), antiprogestogens, antiandrogens (for example flutamide, nilutamide, bicalutamide, cyproterone acetate), LHRH agonists and antagonists (for example goserelin acetate, luprolide), inhibitors of testosterone 5 α -dihydroreductase (for example finasteride), anti-invasion agents (for example metalloproteinase inhibitors like marimastat and inhibitors of urokinase plasminogen activator receptor function) and inhibitors of growth factor function, (such growth factors include for example platelet derived growth factor and hepatocyte growth factor such inhibitors include

growth factor antibodies, growth factor receptor antibodies, tyrosine kinase inhibitors and serine/threonine kinase inhibitors); and

- (iii) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as antimetabolites (for example antifolates like methotrexate, fluoropyrimidines like 5-fluorouracil, purine and adenosine analogues, cytosine arabinoside); antitumour antibiotics (for example anthracyclines like doxorubicin, daunomycin, epirubicin and idarubicin, mitomycin-C, dactinomycin, mithramycin); platinum derivatives (for example cisplatin, carboplatin); alkylating agents (for example nitrogen mustard, melphalan, chlorambucil, busulphan, cyclophosphamide, ifosfamide, nitrosoureas, thiotepa); antimitotic agents (for example vinca alkaloids like vincristine and taxoids like taxol, taxotere); topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan). According to this aspect of the invention there is provided a pharmaceutical product comprising a pyrimidine derivative of the formula (I) as defined hereinbefore, or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, and an additional anti-tumour substance as defined hereinbefore for the conjoint treatment of cancer. An anti-emetic may also be usefully administered, for example when using such conjoint treatment as described above.

In addition to their use in therapeutic medicine, the compounds of formula (I) and their pharmaceutically acceptable salts are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of cell cycle activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

In the above other, pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

The invention will now be illustrated in the following non-limiting Examples, in which standard techniques known to the skilled chemist and techniques analogous to those described in these Examples may be used where appropriate, and in which, unless otherwise stated:

- (i) evaporations were carried out by rotary evaporation in vacuo and work up procedures were carried out after removal of residual solids such as drying agents by filtration;
- (ii) operations were carried out at ambient temperature, typically in the range 18-25°C and in

air unless stated, or unless the skilled person would otherwise operate under an atmosphere of an inert gas such as argon;

(iii) column chromatography (by the flash procedure) and medium pressure liquid chromatography (MPLC) were performed on Merck Kieselgel silica (Art. 9385) or Merck

5 Lichroprep RP-18 (Art. 9303) reversed-phase silica obtained from E. Merck, Darmstadt, Germany;

(iv) yields are given for illustration only and are not necessarily the maximum attainable;

(v) melting points where given were determined using a Mettler SP62 automatic melting point apparatus, an oil-bath apparatus or a Koffler hot plate apparatus.

10 (vi) the structures of the end-products of the formula (I) were generally confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; proton magnetic resonance chemical shift values were measured in deuterated DMSO-d₆ (unless otherwise stated) on the delta scale (ppm downfield from tetramethylsilane) using a Varian Gemini 2000 spectrometer operating at a field strength of 300MHz unless otherwise stated; and peak

15 multiplicities are shown as follows: s, singlet; d, doublet; t, triplet; m, multiplet; br, broad; mass spectrometry (MS) was performed by electrospray on a VG platform;

(vii) intermediates were not generally fully characterised and purity was assessed by thin layer chromatography (TLC), high performance liquid chromatography (HPLC), infra-red (IR), MS or NMR analysis;

20 (viii) the following abbreviations may be used hereinbefore or hereinafter :-

DMF	<i>N,N</i> -dimethylformamide;
NMP	<i>N</i> -methylpyrrolidin-2-one;
DMSO	dimethylsulphoxide;
Rt	retention time;

25 (ix) System A:


Column	4.6mm x 10cm Hichrom RPB 5A
Solvent	A = 95% water, 5% acetonitrile + 0.1% Formic acid B = 95% acetonitrile, 5% water + 0.1% Formic acid
Run time	10 minutes with a 9.5 minute gradient from 5-95% B
30 Wavelength	254nm bandwidth 10nm
Mass detector	Platform LC




Example 1**2-(4-Sulphamoylanilino)-4-(2-cyanoanilino)pyrimidine**


2-Chloro-4-(2-cyanoanilino)pyrimidine (250 mg, 1.09 mmol) was dissolved in *n*-butanol (3 ml) and sulphanilamide (150 mg, 0.87 mmol) was added. The resulting
5 suspension was treated with methanol until all the solid dissolved. The reaction mixture was heated at 95°C for 12 hours and allowed to cool to ambient temperature. The reaction mixture was then basified to pH 9-10 using methanolic ammonia and evaporated onto silica (5 ml). The residue was purified by column chromatography eluting with 0-15% 2.0M methanolic ammonia solution in dichloromethane to afford a solid product (256 mg). NMR (303.1K):
10 6.58 (d, 1H), 7.24 (br s, 2H), 7.56 (m, 5H), 7.69 (d, 1H), 7.82 (t, 1H), 7.95 (d, 1H), 8.16 (d, 1H), 10.88 (br s, 1H), 11.07 (br s, 1H); MS (M+H)⁺: 367.1.


Examples 2-11

The following compounds were prepared by an analogous method to that described in
15 Example 1 using the appropriate 4-sulphonyl aniline and 2-chloro-4-anilinopyrimidine intermediates.



Ex	R ¹	R ²	R ³	R ⁴	NMR, 400MHz @ 373k	MS (M+H) ⁺
2	H	4-Br		H	0.90 (m, 6H), 2.41 (m, 6H), 2.86 (m, 2H), 6.33 (d, 1H), 6.78 (br s, 1H), 7.47 (d, 2H), 7.64 (m, 4H), 7.89 (d, 2H), 8.09 (d, 1H), 9.24 (br s, 2H)	519.3, 521.3
3	-CH ₂ CH ₂ CH ₂ CF ₃	4-Br		H	0.99 (m, 6H), 1.95 (m, 2H), 2.39 (m, 2H), 2.50 (m, 6H), 2.92 (m, 2H), 4.07 (m, 2H), 5.97 (d, 1H), 6.80 (br s, 1H), 7.39 (d, 2H), 7.70 (d, 2H), 7.76 (d, 2H), 7.89 (d, 2H), 8.04 (d, 1H), 9.29 (s, 1H)	629.4, 631.4
4	-CH ₂ CH=CHBr	4-Br		H	0.84 (t, 6H), 2.37 (m, 6H), 2.78 (m, 2H), 4.45 (d, 1H), 4.58 (d, 1H), 5.85 (d, 1H), 6.40 (m, 2H), 6.66 (br s, 1H), 7.24 (d, 2H), 7.56 (d, 2H), 7.61 (d, 2H), 7.75 (dd, 2H), 7.92 (d, 1H), 9.19 (br s, 1H)	637.3, 639.3, 641.3

5	-CH ₂ CH=CHPh	4-Br		H	0.89 (t, 6H), 2.43 (m, 6H), 2.83 (t, 2H), 4.70 (d, 2H), 6.00 (d, 1H), 6.39 (m, 1H), 6.59 (d, 1H), 6.72 (br s, 1H), 7.29 (m, 7H), 7.61 (d, 2H), 7.67 (d, 2H), 7.83 (d, 2H), 8.00 (d, 1H), 9.25 (s, 1H)	635.4, 637.4
6	H	2-CN	-CH ₃	H	3.09 (s, 3H), 6.41 (d, 1H), 7.40 (m, 1H), 7.68 (d, 2H), 7.74 (m, 2H), 7.84 (d, 1H), 7.88 (d, 2H), 8.17 (d, 1H), 9.30 (s, 1H), 9.37 (s, 1H)	366.3
7	H	2-F, 5-CH ₃	-NH ₂	H	(303.1K) 2.28 (s, 3H), 6.55 (d, 1H), 7.16 (m, 1H), 7.25 (m, 3H), 7.46 (d, 1H), 7.64 (t, 4H), 8.09 (d, 1H), 10.71 (br s, 1H), 10.98 (br s, 1H)	374.3
8	-CH ₂ CH ₂ CH ₂ CF ₃	2-F, 5-CH ₃	-CH ₃	H	1.86 (m, 2H), 2.32 (m, 2H), 2.33 (s, 3H), 3.07 (s, 3H), 3.94 (t, 2H), 5.88 (d, 1H), 7.26 (m, 3H), 7.68 (d, 2H), 7.83 (d, 2H), 8.00 (d, 1H), 9.31 (s, 1H)	483.4

9	-CH ₂ CH=CHPh	2-F, 5-CH ₃	-CH ₃	H	2.32 (s, 3H), 3.06 (s, 3H), 4.65 (d, 2H), 5.97 (d, 1H), 6.37 (m, 1H), 6.61 (d, 1H), 7.28 (m, 8H), 7.66 (d, 2H), 7.87 (d, 2H), 8.02 (d, 1H), 9.35 (s, 1H)	489.5
10	-CH ₂ CH=CHBr	2-F, 5-CH ₃	-CH ₃	H	2.33 (s, 3H), 3.08 (s, 3H), 4.48 (d, 1H), 4.61 (d, 1H), 5.89 (m, 1H), 6.40 (m, 1H), 6.54 (m, 1H), 7.24 (m, 3H), 7.69 (d, 2H), 7.85 (d, 2H), 8.01 (d, 1H), 9.37 (s, 1H)	491.3, 493.3
11	-CH ₂ CH=CHPh	2-F, 5-CH ₃	-NH ₂	Cl	2.28 (s, 3H), 4.71 (d, 2H), 6.39 (m, 1H), 6.52 (d, 1H), 6.75 (br s, 2H), 7.10 (m, 2H), 7.19 (m, 2H), 7.27 (m, 4H), 7.69 (d, 2H), 7.80 (d, 2H), 8.08 (s, 1H), 9.38 (br s, 1H) ²	524.5, 526.4
12	H	H		Br	NMR (ambient temp) 2.2 (m, 4H), 2.5 (m, 4H), 3.4 (m, 4H), 7.2 (t, 1H), 7.4 (t, 2H), 7.6 (m, 4H), 7.8 (d, 2H), 8.3 (s, 1H), 8.8 (s, 1H), 9.8 (s, 1H)	518, 520

¹ Product purified by column chromatography twice followed by recrystallization from methanol.

² Run on a 500MHz NMR machine.


5 Example 13

2-(4-Sulphamoylanilino)-4-[2-fluoro-5-methyl-N-(4,4,4-trifluorobutyl)anilino]pyrimidine



2-Chloro-4-(*N*-4,4,4-trifluorobutyl-2-fluoro-5-methylanilino)pyrimidine (215 mg, 0.62 mmol) was dissolved in *n*-butanol (2 ml) and sulphanilamide (85 mg, 0.50 mmol) was added. The resulting suspension was treated with methanol until all the solid dissolved. The reaction mixture was heated at 95°C for 12 hours and allowed to cool to ambient temperature. The solid that had precipitated was collected by filtration, washed with a small volume of methanol and dried in vacuo to yield a white solid (132 mg). NMR (400MHz @ 373K): 1.85 (m, 2H), 2.31 (m, 2H), 2.35 (s, 3H), 3.97 (t, 2H), 6.04 (d, 1H), 7.29 (m, 3H), 7.67 (m, 4H), 8.02 (d, 1H), 10.00 (br s, 1H); MS (M+H)⁺: 484.4.

Examples 14-22

The following compounds were prepared by an analogous method to that described in Example 13 using the appropriate 4-sulphonyl aniline and 2-chloro-4-anilinopyrimidine intermediates.



Ex	R ¹	R ²	R ³	R ⁴	NMR, 400MHz @ 373k	MS (M+H) ⁺
14	-CH ₂ CH=CHBr	2-F, 5-CH ₃	-NH ₂	H	2.34 (s, 3H), 4.50 (d, 1H), 4.66 (d, 1H), 6.00 (m, 1H), 6.39 (m, 1H), 6.58 (m, 1H), 7.29 (m, 3H), 7.69 (m, 4H), 8.05 (d, 1H), 9.99 (br s, 1H)	492.3, 494.3
15	-CH ₂ CH=CHPh	2-F, 5-CH ₃	-NH ₂	H	2.33 (s, 3H), 4.69 (d, 2H), 6.09 (d, 1H), 6.32 (m, 1H), 6.60 (d, 1H), 7.28 (m, 8H), 7.69 (s, 4H), 8.05 (d, 1H), 10.00 (br s, 1H)	490.5
16	-CH ₂ CH ₂ CH ₂ CF ₃	2-CN	-NH ₂	H	1.91 (m, 2H), 2.36 (m, 2H), 4.04 (t, 2H), 6.06 (d, 1H), 7.62 (m, 6H), 7.89 (t, 1H), 7.98 (d, 1H), 8.09 (d, 1H), 9.70 (br s, 1H)	477.4
17	-CH ₂ CH=CHBr	2-CN	-NH ₂	H	4.59 (d, 1H), 4.74 (d, 1H), 6.07 (dd, 1H), 6.48 (m, 1H), 6.65 (m, 1H), 7.65 (m, 6H), 7.90 (m, 1H), 7.99 (d, 1H), 8.12 (d, 1H), 9.79 (br s, 1H)	485.3, 487.3

18	-CH ₂ CH=CHPh	2-CN	-NH ₂	H	(500MHz @ 373K) 4.74 (d, 2H), 6.09 (d, 1H), 6.36 (m, 1H), 6.63 (d, 1H), 7.21 (t, 1H), 7.29 (t, 2H), 7.34 (d, 2H), 7.61 (m, 6H), 7.85 (t, 1H), 7.92 (d, 1H), 8.10 (d, 1H), 9.51 (br s, 1H)	483.4
19	H	H		Br	Rt=2.46 ¹	498
20	H	H	NHCH ₃	Br	Rt=2.56 ¹	435
21	H	H	NH ₂	Br	Rt=2.40 ¹	421
22	H	H		Br	(ambient temperature) 2.7 (s, 3H), 3.38 (m, 2H), 3.6 (m, 2H), 6.55 (d, 2H), 6.6 (t, 1H), 7.1 (d, 2H), 7.2 (t, 1H), 7.4 (t, 2H), 7.6 (m, 4H), 7.8 (d, 2H), 8.3 (s, 1H), 8.8 (br s, 1H), 9.95 (br s, 1H)	538, 540

¹ Chromatography and MS was carried out by LCMS on a Micromass OpenLynx system using System A:

5 **Example 23**


2-(4-Sulphamoylanilino)-4-(2-cyanoanilino)-5-chloropyrimidine

2,5-Dichloro-4-(2-cyanoanilino)pyrimidine (265mg, 1.00 mmol) was dissolved in *n*-butanol (1 ml) and sulphanilamide (207 mg, 1.20 mmol) was added. The resulting suspension was heated at reflux for 2 hours and allowed to cool to ambient temperature. The

reaction mixture was then basified using methanolic ammonia and evaporated onto silica. The residue was purified by column chromatography eluting with 0-15% 2.0M methanolic ammonia solution in dichloromethane to afford a solid product (42.6 mg). NMR (303.1K): 7.09 (2H, s), 7.41-7.59 (5H, m), 7.65 (1H, d), 7.74-7.86 (1H, m), 7.94 (1H, d), 8.25 (1H, s), 9.46 (1H, s), 9.78 (1H, s); MS (M+H)⁺: 401, 403.



Examples 24-28

The following compounds were prepared by an analogous method to that described in Example 23 using the appropriate 4-sulphonyl aniline.



10

Ex	R ⁴	R ⁵	R ²	R ³	NMR, 400MHz @ 373k	MS
24	Cl	H	2-CN		2.46 (3H, s), 7.38-7.58 (6H, m), 7.79 (1H, t), 7.93 (1H, d), 8.25 (1H, s), 9.46 (1H, s), 9.80 (1H, s), 13.79 (1H, s)	497, 499 (M-H)
25	Cl	H	2-CN		3.09 (4H, t), 3.50 (4H, dt), 4.77 (1H, t), 7.46 (2H, d), 7.50 (1H, t), 7.55-7.68 (3H, m), 7.80 (1H, t), 7.94 (1H, d), 8.27 (1H, s), 9.47 (1H, s), 9.83 (1H, s)	487, 489 (M-H)

26	Cl	2-OMe	2-CN	-NH ₂	3.83 (3H, s), 7.08 (3H, m), 7.32-7.41 (1H, m), 7.49 (1H, dd), 7.65-7.71 (2H, m), 7.82 (1H, d), 7.98 (1H, s), 8.22 (1H, s), 8.30-8.35 (1H, m), 9.16 (1H, s)	431, 433 (M+H)
27	Cl	H	2-CN		6.81 (1H, d), 7.22 (1H, d), 7.40-7.57 (5H, m), 7.62 (1H, d), 7.79 (1H, t), 7.91 (1H, d), 8.24 (1H, s), 9.45 (1H, s), 9.76 (1H, s), 12.57 (1H, s)	484, 486 (M+H)
28	Cl	H	2-CN		1.60 (3H, s), 2.08 (3H, s), 7.39 (2H, d), 7.46-7.67 (4H, m), 7.79 (1H, t), 7.91 (1H, d), 8.28 (1H, s), 9.51 (1H, s), 9.91 (1H, s), 10.75 (1H, s)	496, 498 (M+H)

Example 29**2,4-Di-(4-sulphamoylanilino)-5-bromopyrimidine**

- 5 A solution of 5-bromo-2,4-dichloropyrimidine (228 mg, 1.0 mmol), 4-sulphanilamide (180 mg, 1.05 mmol) and *N,N*-diisopropylethylamine (174 μ l, 1.0 mmol) in *n*-butanol (30 ml) was heated at 100°C for 16 hours. A gum formed out of solution. Diethyl ether (20 ml) was added, causing the gum to solidify, and further precipitation to occur. The solid was collected by filtration and triturated with hot methanol, giving the title product (55 mg) as a white solid.
- 10 NMR: (s, 2H), 7.30 (s, 2H), 7.64 (d, 2H), 7.80 (m, 4H), 7.88 (d, 2H), 8.36 (s, 1H), 8.93 (s, 1H), 9.88 (s, 1H).

Example 30**2-(3-Sulphamoylanilino)-4-[(2-morpholino)anilino]pyrimidine**

- 15 In this example the operations were carried out using a Zymate XP robot with solution additions via a Zymate Master Laboratory Station and stirred in a Stem RS5000

Reacto-Station. The structure of the compound was confirmed by LCMS on a Micromass OpenLynx system using the System A.

To 3-aminobenzenesulphonamide (172 mg, 1.0 mmol) in 1,4-dioxane (8 ml) was added 2-Chloro-4-[(2-Morpholino)anilino]pyrimidine (290 mg, 1 mmol) and hydrogen chloride (4.0 M solution in 1,4-dioxane, 50 μ l). The mixture was heated at 100°C for 60 hrs. The reaction mixture was cooled, the resulting solid filtered, washed with 1,4-dioxane and dried (*in vacuo* at 48°C) to give pale brown solid (358 mg). Rt 5.88; MS (M+H)⁺: 427.

Example 31

10 2-(4-Sulphamoylanilino)-4-(4-methoxyphenoxy)-5-chloropyrimidine

A solution of 2,5-dichloro-4-(4-methoxyphenoxy)pyrimidine (Method 25; 0.65 g, 2.4 mmol) and sulphanilamide (0.38 g, 2.2 mmol) in NMP (1.5 ml) was heated to 100°C for 4 hours. The reaction mixture was partitioned between water and ethyl acetate. The organic solution was evaporated and the residue purified by column chromatography eluting with 0.1% formic acid, 4% methanol in dichloromethane to afford a solid product (0.17 g, 19%). NMR: 3.81 (s, 3H), 7.05 (d, 2H), 7.10 (s, 2H), 7.22 (d, 2H), 7.50 (d, 4H), 8.50 (s, 1H), 10.6 (br s, 1H); MS [M-H]⁻: 405, 407.

Preparation of Starting Materials

20 The starting materials for the examples above are either commercially available or are readily prepared by standard methods from known compounds. The following are methods used in the preparation of some of the starting materials used in the above reactions.

Method 1


25 2-Chloro-4-(2-fluoro-5-methyl(N-4,4,4-trifluorobutyl)anilino)pyrimidine

2-Chloro-4-(2-fluoro-5-methylanilino)pyrimidine (750 mg, 3.16 mmol), 4,4,4-trifluoro-1-bromobutane (725 mg, 3.80 mmol) and potassium carbonate (525 mg, 3.80 mmol) were dissolved in *N,N*-Dimethylformamide (3 ml). The reaction mixture was stirred at room temperature for 12 hours and then evaporated onto silica (5 ml) and purified by column chromatography eluting with ethyl acetate (0-40%):isohexane, to yield a solid on evaporation (976 mg). NMR (373K): 1.79 (m, 2H), 2.28 (m, 2H), 2.33 (s, 3H), 3.91 (t, 2H), 6.19 (d, 1H), 7.28 (m, 3H), 8.03 (d, 1H); MS (M+H)⁺: 347, 349.

Methods 2-10

The following compounds were prepared by an analogous method to that described in Method 1 using the appropriate 2-chloro-4-anilinopyrimidine and the relevant alkylating

5 agent.



Method	R ¹	R ²	R ³	MS (M+H) ⁺
2	-CH ₂ CH ₂ CH ₂ CF ₃	4-Br	H	393, 395 ¹
3	-CH ₂ CH=CHBr	4-Br	H	401, 403, 405 ¹
4	-CH ₂ CH=CHPh	4-Br	H	400.2, 402.2
5	-CH ₂ CH=CHBr	2-F, 5-CH ₃	H	355, 357 ¹
6	-CH ₂ CH=CHPh	2-F, 5-CH ₃	H	354.3, 356.3
7	-CH ₂ CH ₂ CH ₂ CF ₃	2-CN	H	340 ¹
8	-CH ₂ CH=CHBr	2-CN	H	348, 350, 352 ¹
9	-CH ₂ CH=CHPh	2-CN	H	347.2, 349.3
10	-CH ₂ CH=CHPh	2-F, 5-CH ₃	Cl	388.3, 390.3

¹where the mass shown is an M⁺.

Method 1110 **2-Chloro-4-(2-cyanoanilino)pyrimidine**


2,4-Dichloropyrimidine (3 g, 0.02 mol), anthranilonitrile (2.38 g, 0.02 mol) and concentrated hydrochloric acid (cat. amount) were added to water (5 ml). The reaction mixture was warmed to 40°C to dissolve the starting materials and the resulting solution was stirred at room temperature for 12 hours. The solid precipitate that had formed was collected
 15 by filtration and dried in vacuo to yield a pale yellow solid (5.14g). NMR: 6.78 (d, 1H), 7.39 (t, 1H), 7.61 (d, 1H), 7.71 (t, 1H), 7.85 (d, 1H), 8.20 (d, 1H), 10.22 (br s, 1H); MS (M+H)⁺: 231.1, 233.1.

Method 12**2-Chloro-4-(4-bromoanilino)pyrimidine**

2,4-Dichloropyrimidine (3g, 20.14 mmol), 4-bromoaniline (3.46g, 20.14 mmol) and di-isopropylethylamine (3.86 ml, 22.15 mmol) were dissolved in *n*-butanol (5 ml). The reaction mixture was heated at 120°C for 12 hours, cooled and evaporated onto silica (5 ml). The residue was purified by column chromatography and eluted with ethyl acetate (50%):isohexane to yield a solid on evaporation (4.77 g). NMR: 6.74 (d, 1H), 7.55 (m, 4H), 8.16 (d, 1H), 10.09 (br s, 1H); MS (M+H)⁺: 284.1, 286.1, 288.0.

10 Methods 13-17

The following compounds were prepared by an analogous method to that described in Method 12 using the appropriate 2,4-dichloropyrimidine and the relevant aniline.



Method	R ²	R ³	NMR, 400MHz	MS (M+H) ⁺
13	2-F, 5-CH ₃	H		238.1, 240.1
14	2-F, 5-CH ₃	Cl	2.29 (s, 3H), 7.17 (m, 3H), 8.35 (s, 1H), 9.49 (s, 1H)	
15	H	Br		282.1, 284.1, 286.1
16	2-CN	Cl		265.1, 267.1
17	2-	H		291, 293

15 Method 18**4-[2-(*N,N*-Diethylamino)ethylamino]benzenesulfonamide**

The above starting material was prepared as described in Therapie, 1965, 20 (4), p. 917-29.

Method 19**2,4,5-Trichloropyrimidine**

5-Chlorouracil (10.0g, 68.5 mmol) was dissolved in phosphorus oxychloride (60 ml) and phosphorus pentachloride (16.0g, 77 mmol) was added. The reaction mixture was then stirred at reflux (110°C) for 16 hrs then allowed to cool to 20°C. The reaction mixture was then poured slowly and carefully into water (200 ml) at 25°C with vigorous stirring. Then stirred well for 90 minutes before addition of EtOAc (250 ml). Organic layer separated off and aqueous layer re-extracted into EtOAc (250 ml). The organic layers were then combined and washed with sodium bicarbonate (200 ml aqueous solution), brine (200 ml) and then evaporated to a yellow liquid. The crude material was purified by column chromatography eluting with dichloromethane to afford the product as a yellow liquid (6.37g, 51%). NMR (CDCl₃): 8.62 (s, 1H); MS (M⁺): 182, 184, 186.

Method 20**15 4-[2-Hydroxy-3-(dimethylamino)propoxy]aniline hydrochloride**

A solution of 4-[2-hydroxy-3-(dimethylamino)propoxy]nitrobenzene (Method 21, 3.75 g) in ethanol (40 ml) was catalytically hydrogenated over 10% palladium-on-carbon (0.4 g) overnight. The catalyst was removed by filtration through diatomaceous earth and the filtrate was concentrated. The residue was dissolved in diethyl ether containing a small amount of isopropanol and ethereal hydrogen chloride (1M, 16 ml) was added. Diethyl ether was removed by evaporation and the solid residue was suspended in isopropanol. The mixture was heated on a steam bath for several minutes and then allowed to cool. The insoluble solid was collected by filtration, washed with isopropanol and ether, and dried to give the product (3.04 g, 72.4%). NMR: 2.80 (s, 6H), 3.15 (m, 2H), 3.88 (m, 2H), 4.25 (m, 1H), 5.93 (br S, 1H), 6.88 (m, 4H); MS (M+H)⁺: 211; C₁₁H₁₈N₂O₂·1.6 HCl requires: C; 49.2, H; 7.4, N; 10.4, Cl; 21.7%; found: C; 49.2, H; 7.2, N; 10.1; Cl; 19.1%.

Method 21**4-[2-Hydroxy-3-(dimethylamino)propoxy]nitrobenzene**

30 4-(2,3-Epoxypropoxy)nitrobenzene (obtained as described in Synthetic Communications, 1994, 24, 833; 4.3 g,) was dissolved in methanol (30 ml) and DMF (10 ml). A solution of dimethylamine in methanol (2M, 17 ml) was added and the mixture was stirred

overnight. Volatile material was removed by evaporation and the residue was partitioned between saturated sodium bicarbonate (100 ml) and ethyl acetate (100 ml). The organic layer was separated and washed with saturated sodium chloride (2 x 100 ml) and dried (MgSO₄). Concentration gave the product as an oil that slowly crystallised under high vacuum (4.79 g, 89.9%). NMR (CDCl₃): 2.33 (s, 6H), 2.98 (m, 1H), 2.54 (m, 1H), 4.00 (m, 3 H), 7.00 (d, 2H), 8.20 (d, 2H); MS (M+H)⁺: 241.

Method 22

4-{2-[(N-Methyl-N-phenyl)amino]ethylsulphonyl}aniline

10 To a solution of 4-{2-[(N-methyl-N-phenyl)amino]ethylsulphonyl}nitrobenzene (Method 23; 10g, 31.25 mmol) in ethanol (100 ml) was added 5 ml of water, 1 ml HCl (conc.) and 25 g of iron pin dust. The reaction was heated at reflux for 3 hours. The reaction was cooled, basified by the addition of caustic and filtered. The iron residue was extracted with a further 100 ml of boiling ethanol and filtered again. To the combined filtrates was added
15 water (400 ml), the product was collected by filtration. The crude material was dissolved in acetone (80 ml) treated with charcoal, filtered and precipitated by the addition of water. The product was collected by filtration. (6g). Mp 158-159°C; NMR: 2.8 (s, 3H), 3.2 (m, 2H), 3.5 (m, 2H), 6.1 (s, 2H), 6.5 (d, 2H), 6.6 (m, 3H), 7.1 (t, 2H), 7.5 (d, 2H); MS (M+H)⁺: 291.

Method 23

4-{2-[(N-Methyl-N-phenyl)amino]ethylsulphonyl}nitrobenzene

To a solution of 1-[(2-chloroethyl)sulphonyl]-4-nitrobenzene (US 5,716,936; 5g, 22.8 mmol) in water (50 ml) was added sodium acetate (3g, 36.6 mmol) and N-methylaniline (3.3 g, 30.8 mmol). The reaction was heated to reflux for 1 hour. After cooling briefly, the hot
25 aqueous supernatant was decanted. The residual oil was washed with cold water. The resulting solid was collected by filtration and air dried. The crude material was recrystallized from benzene/petroleum ether to give red crystals (5.3g). Mp 111-113°C. This material was used without further characterisation.

Method 24**4-[(2-[N-Morpholino]ethyl)sulphonyl]aniline**

The title compound can be prepared by a method analogous to that used in Method 22 above (parts A-C) using morpholine in place of *N*-methylaniline. NMR: 2.2 (m, 4H), 3.3 (m, 4H), 3.4 (m, 4H), 6.05 (br s, 2H), 6.6 (d, 2H), 7.4 (d, 2H); MS (M+H)⁺: 271.

Method 25**2,5-Dichloro-4-(4-methoxyphenoxy)pyrimidine**

10 A solution of 2,4,5-trichloropyrimidine (1.0 g, 5.4 mmol) and 4-methoxyphenol (0.64 g, 5.2 mmol) in NMP (2.5 ml) was treated with anhydrous potassium carbonate (1.65 g, 12 mmol). The reaction mixture was allowed to stir at ambient temperature for 18 hours. The reaction was partitioned between water and ethyl acetate. The organic layer was evaporated and the residue purified by column chromatography eluting with 10% ethyl acetate in
15 isohexane to afford a solid product (1.33 g, 90%). NMR: 3.78 (s, 3H), 7.00 (d, 2H), 7.20 (d, 2H), 8.77 (s, 1H); MS (M+H)⁺ 271, 273.

Example 32

The following illustrate representative pharmaceutical dosage forms containing the
20 compound of formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof (hereafter compound X), for therapeutic or prophylactic use in humans:-

(a): Tablet I	mg/tablet
Compound X	100
Lactose Ph.Eur	182.75
Croscarmellose sodium	12.0
Maize starch paste (5% w/v paste)	2.25
Magnesium stearate	3.0

(b): Tablet II	mg/tablet
Compound X	50
Lactose Ph.Eur	223.75
Croscarmellose sodium	6.0
Maize starch	15.0
Polyvinylpyrrolidone (5% w/v paste)	2.25
Magnesium stearate	3.0

(c): Tablet III	mg/tablet
Compound X	1.0
Lactose Ph.Eur	93.25
Croscarmellose sodium	4.0
Maize starch paste (5% w/v paste)	0.75
Magnesium stearate	1.0

(d): Capsule	mg/capsule
Compound X	10
Lactose Ph.Eur	488.5
Magnesium stearate	1.5

(e): Injection I	(50 mg/ml)
Compound X	5.0% w/v
1M Sodium hydroxide solution	15.0% v/v
0.1M Hydrochloric acid	(to adjust pH to 7.6)
Polyethylene glycol 400	4.5% w/v
Water for injection	to 100%

(f): Injection II	10 mg/ml
Compound X	1.0% w/v
Sodium phosphate BP	3.6% w/v
0.1M Sodium hydroxide solution	15.0% v/v
Water for injection	to 100%


(g): Injection III	(1mg/ml, buffered to pH6)
Compound X	0.1% w/v
Sodium phosphate BP	2.26% w/v
Citric acid	0.38% w/v
Polyethylene glycol 400	3.5% w/v
Water for injection	to 100%

Note

- 5 The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example to provide a coating of cellulose acetate phthalate.

Claims

1. A pyrimidine derivative of the formula (I):




5


(I)

wherein:

- Q_1 and Q_2 are independently selected from aryl or carbon linked heteroaryl; and one of Q_1 and Q_2 or both Q_1 and Q_2 is substituted on a ring carbon by one group selected from sulphamoyl, *N*-(C_{1-4} alkyl)sulphamoyl (optionally substituted by halo or hydroxy),
- 10 *N,N*-di-(C_{1-4} alkyl)sulphamoyl (optionally substituted by halo or hydroxy), C_{1-4} alkylsulphonyl (optionally substituted by halo or hydroxy) or a substituent of the formula (Ia) or (Ia'):



(Ia)



(Ia')

wherein:

- 15 Y is -NHS(O)₂-, -S(O)₂NH- or -S(O)₂-;

Z is R^aO -, R^bR^cN -, R^dS -, $R^eR^fNNR^g$ -, C_{3-8} cycloalkyl, phenyl or a heterocyclic group; wherein said phenyl, C_{3-8} cycloalkyl or heterocyclic group are optionally substituted on a ring carbon by one or more groups selected from R^h ; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from

- 20 R^i ;

R^a , R^b , R^c , R^d , R^e , R^f and R^g are independently selected from hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, phenyl, heterocyclic group and C_{3-8} cycloalkyl; wherein said C_{1-4} alkyl, C_{2-4} alkenyl and C_{3-8} cycloalkyl are optionally substituted by one or more groups selected from R^j ;

n is 0 or 1;

m is 1, 2 or 3, in addition **m** may be 0 when **Z** is C₃₋₈cycloalkyl, phenyl or a heterocyclic group;

Q₃ is a nitrogen linked heterocycle; wherein said heterocycle is optionally substituted on a ring carbon by one or more groups selected from **R^k**; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from **R^m**

G is -O-, -S- or -NR²-;

R² is selected from hydrogen, C₁₋₆alkyl, C₃₋₆alkenyl and C₃₋₆alkynyl; wherein said C₁₋₆alkyl, C₃₋₆alkenyl and C₃₋₆alkynyl are optionally substituted by one or more groups selected from **Rⁿ**;

R¹ is selected from hydrogen, halo, hydroxy, nitro, amino, *N*-(C₁₋₃alkyl)amino, *N,N*-di-(C₁₋₃alkyl)amino, cyano, trifluoromethyl, trichloromethyl, C₁₋₃alkyl [optionally substituted by 1 or 2 substituents independently selected from halo, cyano, amino, *N*-(C₁₋₃alkyl)amino, *N,N*-di-(C₁₋₃alkyl)amino, hydroxy and trifluoromethyl], C₃₋₅alkenyl [optionally substituted by up to three halo substituents, or by one trifluoromethyl substituent], C₃₋₅alkynyl, C₁₋₃alkoxy, mercapto, C₁₋₃alkylsulphanyl, carboxy and C₁₋₃alkoxycarbonyl;

Q₁ is optionally substituted on a ring carbon by one to four substituents independently selected from halo, mercapto, nitro, formyl, formamido, carboxy, cyano, amino, ureido, carbamoyl, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl [wherein said C₁₋₄alkyl, C₂₋₄alkenyl and C₂₋₄alkynyl are optionally substituted by one or more groups selected from **R^o**], C₁₋₄alkanoyl, C₁₋₄alkoxycarbonyl, heterocyclic group, C₁₋₄alkylS(O)_a wherein **a** is 0 or 1 [optionally substituted by hydroxy], *N'*-(C₁₋₄alkyl)ureido, *N',N'*-di-(C₁₋₄alkyl)ureido, *N*-(C₁₋₄alkyl)-*N*-(C₁₋₄alkyl)ureido, *N',N'*-di-(C₁₋₄alkyl)-*N*-(C₁₋₄alkyl)ureido, *N*-C₁₋₄alkylamino, *N,N*-di-(C₁₋₄alkyl)amino, *N*-C₁₋₄alkylcarbamoyl, *N,N*-di-(C₁₋₄alkyl)carbamoyl and C₁₋₄alkanoylamino;

and also independently, or in addition to, the above substituents, **Q₁** may be optionally substituted by one to two substituents independently selected from aryl, C₃₋₈cycloalkyl and a heterocyclic group; wherein said aryl, C₃₋₈cycloalkyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from **R^p**; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from **R^q**;

and also independently, or in addition to, the above substituents, Q_1 may be optionally substituted by one C_{1-4} alkoxy or by one hydroxy substituent;

Q_2 is optionally substituted on a ring carbon by one to four substituents independently selected from halo, hydroxy, mercapto, nitro, formyl, formamido, carboxy, cyano, amino, ureido, carbamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy [wherein said C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl and C_{1-4} alkoxy are optionally substituted by one or more groups selected from R^7], C_{1-4} alkanoyl, C_{1-4} alkoxycarbonyl, heterocyclic group, C_{1-4} alkylS(O)_a, wherein a is 0 or 1 [optionally substituted by hydroxy], $N'-(C_{1-4}$ alkyl)ureido, N',N' -di-(C_{1-4} alkyl)ureido, $N'-(C_{1-4}$ alkyl)- N -(C_{1-4} alkyl)ureido, N',N' -di-(C_{1-4} alkyl)- N -(C_{1-4} alkyl)ureido, N - C_{1-4} alkylamino, N,N -di-(C_{1-4} alkyl)amino, N - C_{1-4} alkylcarbamoyl, N,N -di-(C_{1-4} alkyl)carbamoyl, C_{2-4} alkenyloxy, C_{2-4} alkynyloxy and C_{1-4} alkanoylamino;

and also independently, or in addition to, the above substituents, Q_2 may be optionally substituted by one to two substituents independently selected from aryl, C_{3-8} cycloalkyl or a heterocyclic group; wherein said aryl, C_{3-8} cycloalkyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R^5 ; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^4 ;

R^j , R^n , R^o and R^r are independently selected from hydroxy, halo, amino, cyano, formyl, formamido, carboxy, nitro, mercapto, carbamoyl, sulphamoyl, N - C_{1-4} alkylamino, N,N -di-(C_{1-4} alkyl)amino, C_{1-4} alkanoyl, C_{1-4} alkanoyloxy, C_{1-4} alkoxy, C_{1-4} alkoxycarbonyl, N - C_{1-4} alkylcarbamoyl, N,N -di-(C_{1-4} alkyl)carbamoyl, C_{1-4} alkanoylamino, C_{1-4} alkylS(O)_a, wherein a is 0 to 2, C_{1-4} alkylsulphonylamino, N -(C_{1-4} alkyl)sulphamoyl, N -(C_{1-4} alkyl)₂sulphamoyl, N -(C_{1-4} alkyl)carbamoyl, N -(C_{1-4} alkyl)₂carbamoyl, phenyl, phenylthio, phenoxy, C_{3-8} cycloalkyl and a heterocyclic group; wherein said phenyl, phenylthio, phenoxy, C_{3-8} cycloalkyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R^u ; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^v ;

R^h , R^k , R^p , R^s and R^u are independently selected from hydroxy, halo, amino, cyano, formyl, formamido, carboxy, nitro, mercapto, carbamoyl, sulphamoyl, C_{1-4} alkyl [optionally substituted by one or more groups selected from halo, cyano, amino, N - C_{1-4} alkylamino,

N,N-di-(C₁₋₄alkyl)amino or hydroxy], C₂₋₄alkenyl [optionally substituted by one or more groups selected from halo], C₂₋₄alkynyl, *N*-C₁₋₄alkylamino, *N,N*-di-(C₁₋₄alkyl)amino, C₁₋₄alkanoyl, C₁₋₄alkanoyloxy, C₁₋₄alkoxy [optionally substituted by one or more groups selected from halo], C₁₋₄alkoxycarbonyl, *N*-C₁₋₄alkylcarbamoyl, *N,N*-di-(C₁₋₄alkyl)carbamoyl, 5 C₁₋₄alkanoylamino, C₁₋₄alkylS(O)_a wherein a is 0 to 2, C₁₋₄alkylsulphonylamino, *N*-(C₁₋₄alkyl)sulphamoyl, *N*-(C₁₋₄alkyl)₂sulphamoyl, phenyl, C₃₋₈cycloalkyl and a heterocyclic group; and

Rⁱ, R^q, R^t and R^v are independently selected from C₁₋₄alkyl, C₁₋₄alkanoyl, C₁₋₄alkylsulphonyl, C₁₋₄alkoxycarbonyl, carbamoyl, *N*-(C₁₋₄alkyl)carbamoyl, 10 *N,N*-(C₁₋₄alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl; or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

2. A pyrimidine derivative as claimed in claim 1 wherein Q₁ is phenyl or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

15

3. A pyrimidine derivative as claimed in either of claims 1 or 2 wherein Q₂ is phenyl or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

4. A pyrimidine derivative as claimed in any one of claims 1 to 3 wherein one of Q₁ and 20 Q₂ or both of Q₁ and Q₂ are substituted on a ring carbon by one group selected from sulphamoyl, mesyl, *N*-(2-diethylaminoethyl)sulphamoyl, 2-(*N*-methyl-*N*-phenylamino)ethylsulphonyl, 2-morpholinoethylsulphonyl, *N*-(5-methylthiadiazol-2-yl)sulphamoyl, *N,N*-di-(2-hydroxyethyl)sulphamoyl, *N*-(thiazol-2-yl)sulphamoyl, *N*-(3,4-dimethylisoxazol-5-yl)sulphamoyl, 25 *N*-(pyrid-2-yl)sulphamoyl and *N*-methylsulphamoyl or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

5. A pyrimidine derivative as claimed in any one of claims 1 to 3 wherein Q₁ is phenyl substituted in the para- or meta- position relative to the -NH- by sulphamoyl, 30 *N*-(C₁₋₄alkyl)sulphamoyl (optionally substituted by halo or hydroxy), *N,N*-di-(C₁₋₄alkyl)sulphamoyl (optionally substituted by halo or hydroxy), C₁₋₄alkylsulphonyl

(optionally substituted by halo or hydroxy) or a substituent of the formula (Ia) or (Ia') or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

6. A pyrimidine derivative as claimed in any one of claims 1 to 5 wherein G is -O-, -NH-,
5 -(4,4,4-trifluorobutyl)N-, -(3-bromo-2-propenyl)N- or -(3-phenyl-2-propenyl)N- or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

7. A pyrimidine derivative as claimed in any one of claims 1 to 6 wherein R¹ is hydrogen or halo or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

10

8. A pyrimidine derivative as claimed in any one of claims 1 to 7 wherein Q₁ is optionally substituted by one C₁₋₄alkoxy substituent or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

15 9. A pyrimidine derivative as claimed in any one of claims 1 to 8 wherein Q₂ is optionally substituted on a ring carbon by one to two substituents independently selected from halo, cyano, methyl, methoxy and morpholino or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

20 10. A pyrimidine derivative as claimed in any one of claims 1 to 9 selected from:

2-(4-sulphamoylanilino)-4-(2-cyanoanilino)pyrimidine;

2-(4-N-methylsulphamoylanilino)-4-anilino-5-bromopyrimidine;


2-(4-sulphamoylanilino)-4-anilino-5-bromopyrimidine;

2,4-di-(4-sulphamoylanilino)-5-bromopyrimidine; or

25 2-(4-sulphamoylanilino)-4-(4-methoxyphenoxy)-5-chloropyrimidine;
or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.


11. A process for preparing a pyrimidine derivative, or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1 to 10 which comprises
30 of:-

a) for compounds of formula (I) where G is -NR²-; reacting a pyrimidine of formula (II):



(II)

wherein L is a displaceable group as defined below, with a compound of formula (III):




(III)

5


where G is -NR²-;

b) reaction of a pyrimidine of formula (IV):




(IV)

10 wherein L is a displaceable group as defined below, with a compound of formula (V):



(V)


c) for compounds of formula (I) wherein the sidechain is of formula (Ia) and Y is -S(O)₂NH-;
by reaction of a compound of formula (VI):



(VI)


15

where L is a displaceable group; with an amine of formula (VII):




(VII)

d) for compounds of formula (I) wherein the sidechain is of formula (Ia) and Y is -NHS(O)₂- by reaction of an amine of formula (VIII):




5 with a compound of formula (IX):



where L is a displaceable group;

e) for compounds of formula (I) wherein the sidechain is of formula (Ia'); by reaction of a
10 compound of formula (VI) with an amine of formula (X):



and thereafter if necessary:

- i) converting a compound of the formula (I) into another compound of the formula (I);
- 15 ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

12. A pharmaceutical composition which comprises a pyrimidine derivative of formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in
20 any one of claims 1 to 10, in association with a pharmaceutically acceptable diluent or carrier.

13. A pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1 to 10, for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man.

25

14. A pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1 to 10, for use as a medicament.

15. The use of a pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1 to 10, in the manufacture of a medicament for use in the production of an anti-cancer, cell cycle inhibitory
5 (anti-cell-proliferation) effect in a warm-blooded animal such as man.

16. According to a further feature of this aspect of the invention there is provided a method for producing an anti-cancer, cell cycle inhibitory (anti-cell-proliferation) effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering
10 to said animal an effective amount of a pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1 to 10.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 01/00782

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/48 C07D401/12 C07D417/12 C07D413/12 A61K31/505
A61K31/506

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, BEILSTEIN Data, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EL-KERDAWY, M. M. ET AL: "2,4-bis(substituted) 5-nitropyrimidines of expected diuretic action" EGYPT. J. CHEM. (1987), VOLUME DATE 1986, 29(2), 247-51 , XP000999986 see Table 1	1-6, 11-14
X	GHOSH, DOLLY: "2,4-Bis(arylamino)-6-methylpyrimidines as antimicrobial agents" J. INDIAN CHEM. SOC. (1981), 58(5), 512-13 , XP000918018 see Table 1	1-6, 11-14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

5 June 2001

Date of mailing of the international search report

13/06/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Scruton-Evans, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/00782

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 00 12485 A (BREault GLORIA ANNE ; PEASE JANET ELIZABETH (GB); ZENeca LTD (GB)) 9 March 2000 (2000-03-09)	1-16
P, Y	see definitions of formula Ia (especially Z) and optional substituents for Q1 (eg alkylsulphonyl) the whole document	1-16
P, X	WO 00 39101 A (BREault GLORIA ANNE ; PEASE JANET ELIZABETH (GB); ASTRAZENECA UK LT) 6 July 2000 (2000-07-06)	1-16
P, Y	see definitions of Ia (especially Z) and optional substituents for Q1 eg alkylsulphonyl the whole document	1-16
P, A	WO 00 59892 A (BREault GLORIA ANNE ; JAMES STEWART RUSSELL (GB); PEASE JANET ELIZA) 12 October 2000 (2000-10-12) the whole document	1-16
A	EP 0 945 443 A (JANSSEN PHARMACEUTICA NV) 29 September 1999 (1999-09-29) the whole document	1-16
Y	WO 97 19065 A (CELLTECH THERAPEUTICS LTD ; DAVIS PETER DAVID (GB); MOFFAT DAVID FE) 29 May 1997 (1997-05-29) see definitions of R7 and its possible substituents (page 6, lines 20ff)	1-16

INTERNATIONAL SEARCH REPORT

International application No.

PCT/ GB/01 / 00782

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

SEE ANNEXE

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-9(partly),11-16(partly)

Present claims 1-9 and 11-16 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds of the examples and claim 1 wherein Q1 is limited to phenyl substituted by an SO₂- group in any position.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/00782

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0012485	A	09-03-2000	AU 5438299 A	21-03-2000
WO 0039101	A	06-07-2000	AU 1874300 A	31-07-2000
WO 0059892	A	12-10-2000	AU 3569800 A	23-10-2000
EP 0945443	A	29-09-1999	AU 3599699 A	18-10-1999
			BR 9909191 A	05-12-2000
			WO 9950250 A	07-10-1999
			NO 20004810 A	26-09-2000
			TR 200002760 T	21-12-2000
			US 6197779 B	06-03-2001
			EP 0945442 A	29-09-1999
WO 9719065	A	29-05-1997	AU 7631496 A	11-06-1997
			EP 0862560 A	09-09-1998
			US 5958935 A	28-09-1999